

# Mathematics for Informatics

## Numerical Mathematics 2 (lecture 9 of 12)

Francesco Dolce

`francesco.dolce@fjfi.cvut.cz`

Czech Technical University in Prague

Fall 2019/2020

created: December 9, 2019, 18:09

# Outline

- 1 Eigenvectors
- 2 Power method
- 3 QR algoritmus

# Outline

- 1 Eigenvectors
- 2 Power method
- 3 QR algoritmus

# Eigenvalues and eigenvectors

A complex number  $\lambda$  is called an **eigenvalue** of the matrix  $M \in \mathbb{C}^{n,n}$ , whenever there exists a non-zero vector  $u \in \mathbb{C}^n$  such that

$$Mu = \lambda u.$$

The vector  $u$  is called an **eigenvector** of the matrix  $M$  relative to the eigenvalue  $\lambda$ .

# Eigenvalues and eigenvectors

A complex number  $\lambda$  is called an **eigenvalue** of the matrix  $M \in \mathbb{C}^{n,n}$ , whenever there exists a non-zero vector  $u \in \mathbb{C}^n$  such that

$$Mu = \lambda u.$$

The vector  $u$  is called an **eigenvector** of the matrix  $M$  relative to the eigenvalue  $\lambda$ .

The set of eigenvectors of  $M$  (relative to the eigenvalues  $\lambda$  and to the zero vector) form a base of the subspace  $\ker(M - \lambda E)$ .

# Eigenvalues and eigenvectors

A complex number  $\lambda$  is called an **eigenvalue** of the matrix  $M \in \mathbb{C}^{n,n}$ , whenever there exists a non-zero vector  $u \in \mathbb{C}^n$  such that

$$Mu = \lambda u.$$

The vector  $u$  is called an **eigenvector** of the matrix  $M$  relative to the eigenvalue  $\lambda$ .

The set of eigenvectors of  $M$  (relative to the eigenvalues  $\lambda$  and to the zero vector) form a base of the subspace  $\ker(M - \lambda E)$ .

The eigenvalues of the matrix  $M$  are the roots of the **characteristic polynomial** of the  $M$ , that is the polynomial

$$p_M(\lambda) := \det(M - \lambda E).$$

Therefore, each matrix  $M \in \mathbb{C}^{n,n}$  has at most  $n$  different complex eigenvalues.

# Diagonalizability of a matrix

A matrix  $M \in \mathbb{C}^{n,n}$  is **diagonalizable** when there exist a diagonal matrix  $D \in \mathbb{C}^{n,n}$  and a regular matrix  $P \in \mathbb{C}^{n,n}$  such that

$$M = PDP^{-1}.$$

where  $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ .

# Diagonalizability of a matrix

A matrix  $M \in \mathbb{C}^{n,n}$  is **diagonalizable** when there exist a diagonal matrix  $D \in \mathbb{C}^{n,n}$  and a regular matrix  $P \in \mathbb{C}^{n,n}$  such that

$$M = PDP^{-1}.$$

where  $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ .

**Remind:** In the previous lecture we saw that  $M^k = PD^kP^{-1}$ .



# Diagonalizability of a matrix

A matrix  $M \in \mathbb{C}^{n,n}$  is **diagonalizable** when there exist a diagonal matrix  $D \in \mathbb{C}^{n,n}$  and a regular matrix  $P \in \mathbb{C}^{n,n}$  such that

$$M = PDP^{-1}.$$

where  $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ .

**Remind:** In the previous lecture we saw that  $M^k = PD^kP^{-1}$ .

**Remark:** The columns of the matrix  $P$  are the eigenvectors of  $M$ . These eigenvectors form a basis of  $\mathbb{C}^n$ . The elements of the diagonal matrix  $D$  are the eigenvalues of  $M$  (with their multiplicity).

# Looking for an eigenvector

Let  $M \in \mathbb{C}^{n,n}$ . Suppose it is diagonalizable and we can order its eigenvalues as follows

$$|\lambda_1| > |\lambda_2| \geq \dots \geq |\lambda_n|.$$

# Looking for an eigenvector

Let  $M \in \mathbb{C}^{n,n}$ . Suppose it is diagonalizable and we can order its eigenvalues as follows

$$|\lambda_1| > |\lambda_2| \geq \dots \geq |\lambda_n|.$$

We are looking for the eigenvector of the eigenvalue  $\lambda_1$ , the so-called **dominant eigenvalue**. It is a vector  $u_1$  such that

$$Mu_1 = \lambda_1 u_1.$$

# Looking for an eigenvector

Let  $M \in \mathbb{C}^{n,n}$ . Suppose it is diagonalizable and we can order its eigenvalues as follows

$$|\lambda_1| > |\lambda_2| \geq \dots \geq |\lambda_n|.$$

We are looking for the eigenvector of the eigenvalue  $\lambda_1$ , the so-called **dominant eigenvalue**. It is a vector  $u_1$  such that

$$Mu_1 = \lambda_1 u_1.$$

In general, the matrix need not be diagonalizable, but the ideas would be more complicated (actually, we only require to have one eigenvalue which is the greatest in absolute value).

# Applications

Eigenvalues play an important role in several applications:

- Classification of conics and quadratic forms (geometry).
- Quantum computation, quantum mechanics, asymptotic behaviour of dynamical systems (physics).
- PCA, or *Principal Component Analysis* (big data).
- Recognition of 2D and 3D objects using spectral methods (AI).

# Applications

Eigenvalues play an important role in several applications:

- Classification of conics and quadratic forms (geometry).
- Quantum computation, quantum mechanics, asymptotic behaviour of dynamical systems (physics).
- PCA, or *Principal Component Analysis* (big data).
- Recognition of 2D and 3D objects using spectral methods (AI).
- More practical example: **PageRank** measures a relative importance of WWW documents by inspecting links between them.
  - Its value is in fact an eigenvector of the dominant eigenvalues of a modified adjacency matrix of these links. This matrix satisfies requirements of our problem.
  - **PageRank** is calculated using **power methods**.

# Outline

- 1 Eigenvectors
- 2 Power method
- 3 QR algoritmus

# Introduction and assumptions (1/2)

In its basic variant, the power method is used to find the dominant eigenvalue of a matrix,



# Introduction and assumptions (1/2)

In its basic variant, the power method is used to find the dominant eigenvalue of a matrix,

Given a matrix  $M \in \mathbb{C}^{n,n}$  let us consider a regular matrix  $P \in \mathbb{C}^{n,n}$  such that

$$M = PDP^{-1}$$

where  $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ . Let also suppose that the values are ordered:

$$|\lambda_1| > |\lambda_2| \geq \dots \geq |\lambda_n|.$$

# Introduction and assumptions (1/2)

In its basic variant, the power method is used to find the dominant eigenvalue of a matrix,

Given a matrix  $M \in \mathbb{C}^{n,n}$  let us consider a regular matrix  $P \in \mathbb{C}^{n,n}$  such that

$$M = PDP^{-1}$$

where  $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ . Let also suppose that the values are ordered:

$$|\lambda_1| > |\lambda_2| \geq \dots \geq |\lambda_n|.$$

**Note:** We suppose that the dominant eigenvalue  $\lambda_1$  is not degenerate (i.e., that the corresponding eigenspace has dimension 1).

# Introduction and assumptions (2/2)

We are looking for an eigenvector associated to the eigenvalue  $\lambda_1$ , that is a non-zero vector  $u_1$  such that

$$Mu_1 = \lambda_1 u_1.$$

# Introduction and assumptions (2/2)

We are looking for an eigenvector associated to the eigenvalue  $\lambda_1$ , that is a non-zero vector  $u_1$  such that

$$Mu_1 = \lambda_1 u_1.$$

The **power method** is an **iterative method**. We will construct a sequence  $(x_k)$  as follows:  $x_0$  is chosen randomly and the next terms are determined by

$$x_k = Mx_{k-1} \quad \text{for } k > 0.$$

Equivalently, we have

$$x_k = M^k x_0 \quad k \in \mathbb{N}_0.$$

# Power method principle (1/4)

If  $M$  is regular, thus diagonalizable, there exist eigenvectors  $\{u_1, u_2, \dots, u_n\}$ , which form a basis of  $\mathbb{C}^{n,1}$ .

If  $M$  is not regular, then we need to complete the set of eigenvectors by a basis of the kernel of  $M$ .

# Power method principle (1/4)

If  $M$  is regular, thus diagonalizable, there exist eigenvectors  $\{u_1, u_2, \dots, u_n\}$ , which form a basis of  $\mathbb{C}^{n,1}$ .

If  $M$  is not regular, then we need to complete the set of eigenvectors by a basis of the kernel of  $M$ .

The vector  $x_0$  can be written as  $x_0 = \alpha_1 u_1 + \dots + \alpha_n u_n$ .

Suppose that  $\alpha_1 \neq 0$ .

# Power method principle (1/4)

If  $M$  is regular, thus diagonalizable, there exist eigenvectors  $\{u_1, u_2, \dots, u_n\}$ , which form a basis of  $\mathbb{C}^{n,1}$ .

If  $M$  is not regular, then we need to complete the set of eigenvectors by a basis of the kernel of  $M$ .

The vector  $x_0$  can be written as  $x_0 = \alpha_1 u_1 + \dots + \alpha_n u_n$ .

Suppose that  $\alpha_1 \neq 0$ .

Coefficients  $\alpha_i$  can be absorbed by the eigenvectors ( $u'_i = \alpha_i u_i$ ) and we have

$$x_0 = u'_1 + \dots + u'_n.$$

# Power method principle (2/4)

The recurrent definition of  $x_k$  implies

$$\begin{aligned}x_k &= M^k x_0 \\ &= M^k u_1 + \cdots + M^k u_n \\ &= \lambda_1^k u_1 + \cdots + \lambda_n^k u_n.\end{aligned}$$



# Power method principle (2/4)

The recurrent definition of  $x_k$  implies

$$\begin{aligned}x_k &= M^k x_0 \\ &= M^k u_1 + \cdots + M^k u_n \\ &= \lambda_1^k u_1 + \cdots + \lambda_n^k u_n.\end{aligned}$$

The last equality gives

$$x_k = \lambda_1^k \left( u_1 + \left( \frac{\lambda_2}{\lambda_1} \right)^k u_2 + \cdots + \left( \frac{\lambda_n}{\lambda_1} \right)^k u_n \right).$$

# Power method principle (2/4)

The recurrent definition of  $x_k$  implies

$$\begin{aligned} x_k &= M^k x_0 \\ &= M^k u_1 + \cdots + M^k u_n \\ &= \lambda_1^k u_1 + \cdots + \lambda_n^k u_n. \end{aligned}$$

The last equality gives

$$x_k = \lambda_1^k \left( u_1 + \left( \frac{\lambda_2}{\lambda_1} \right)^k u_2 + \cdots + \left( \frac{\lambda_n}{\lambda_1} \right)^k u_n \right).$$

We rewrite it as

$$x_k = \lambda_1^k (u_1 + \varepsilon_k).$$

Since for all  $j > 1$  we have  $\left| \frac{\lambda_j}{\lambda_1} \right| < 1$ , then  $\lim_{k \rightarrow +\infty} \varepsilon_k = 0$ .

# Power method principle (3/4)

The sequence  $\left( \frac{x_k}{\lambda_1^k} \right)$  “converges” to the eigenvector of the dominant eigenvalues.

# Power method principle (3/4)

The sequence  $\left(\frac{x_k}{\lambda_1^k}\right)$  “converges” to the eigenvector of the dominant eigenvalues.

We have  $\|x_k\| \rightarrow +\infty$ . Thus we need to control the norm: we may set it to **1** at each step (by *normalizing*, i.e., considering  $y_k = \frac{x_k}{\|x_k\|}$ ).

To have convergence also for the case  $\lambda_1 < 0$ , we need to pick the right direction for the eigenvector so that it does not oscillate. We may do this by setting the largest entry in absolute value to **1** (and thus use the maximum norm).

# Power method principle (3/4)

The sequence  $\begin{pmatrix} x_k \\ \lambda_1^k \end{pmatrix}$  “converges” to the eigenvector of the dominant eigenvalues.

We have  $\|x_k\| \rightarrow +\infty$ . Thus we need to control the norm: we may set it to **1** at each step (by *normalizing*, i.e., considering  $y_k = \frac{x_k}{\|x_k\|}$ ).

To have convergence also for the case  $\lambda_1 < 0$ , we need to pick the right direction for the eigenvector so that it does not oscillate. We may do this by setting the largest entry in absolute value to **1** (and thus use the maximum norm).

The speed of convergence is given by  $\lambda_2$  since  $\|\varepsilon_k\| = \mathcal{O}\left(\left|\frac{\lambda_2}{\lambda_1}\right|^k\right)$

# Power method principle (4/4)

How to find the dominant eigenvalue?

If  $\varphi$  is a linear mapping  $\varphi : \mathbb{C}^{n,1} \mapsto \mathbb{C}$  such that  $\varphi(u_1) \neq 0$ , then

$$\frac{\varphi(x_{k+1})}{\varphi(x_k)} = \frac{\varphi(\lambda_1^{k+1}(u_1 + \varepsilon_{k+1}))}{\varphi(\lambda_1^k(u_1 + \varepsilon_k))} = \frac{\lambda_1^{k+1}(\varphi(u_1) + \varphi(\varepsilon_{k+1}))}{\lambda_1^k(\varphi(u_1) + \varphi(\varepsilon_k))} \rightarrow \lambda_1 \quad \text{for } k \rightarrow +\infty.$$

The mapping  $\varphi$  can be set to the mapping defined for all  $x \in \mathbb{C}^{n,1}$  as  $\varphi(x) = x_{(1)}$  where  $x_{(1)}$  is the first component  $x$  (if  $\varphi(u_1) \neq 0$ ).

Power method - demonstration in  $\mathbb{R}^{n,n}$ 

Let us find the dominant eigenvector of  $M = \begin{pmatrix} 2 & 1 \\ 1 & 4 \end{pmatrix}$ , which satisfies the conditions of power method.

The exact solution is  $u_1 = (1, \sqrt{2} + 1) = \frac{1}{\sqrt{2} + 1}(\sqrt{2} - 1, 1)$ , with eigenvalue  $\lambda_1 = 3 + \sqrt{2}$ .

$k$	$\hat{x}_k$	$\ \hat{x}_k - \hat{x}_{k-1}\ _\infty$
0	(1.0, 1.0)	-
1	(0.59999999999999998, 1.0)	0.4
2	(0.47826086956521746, 1.0)	0.121739130435
3	(0.43689320388349517, 1.0)	0.0413676656817
4	(0.42231947483588622, 1.0)	0.0145737290476
5	(0.4171202375061851, 1.0)	0.0051992373297

In the calculations, the maximum entry in absolute value is set to 1 at each step and the convergence criterion  $\|\hat{x}_k - \hat{x}_{k-1}\|_\infty < 10^{-2}$ .

Power method - demonstration in  $\mathbb{C}^{n,n}$  (1/2)

Let us consider the matrix

$$M = \begin{pmatrix} 36408 + 16769i & -5412 - 2481i & 107256 + 49397i & -492 - 214i \\ -10656 - 5164i & 1584 + 762i & -31392 - 15210i & 144 + 66i \\ -12876 - 5954i & 1914 + 881i & -37932 - 17539i & 174 + 76i \\ 4329 - 262i & -643 + 39i & 12753 - 771i & -58 + 6i \end{pmatrix}$$

The eigenvalues are  $-2i$ ,  $-i$ ,  $3i/2$  and  $3/2$ .



Power method - demonstration in  $\mathbb{C}^{n,n}$  (1/2)


Let us consider the matrix

$$M = \begin{pmatrix} 36408 + 16769i & -5412 - 2481i & 107256 + 49397i & -492 - 214i \\ -10656 - 5164i & 1584 + 762i & -31392 - 15210i & 144 + 66i \\ -12876 - 5954i & 1914 + 881i & -37932 - 17539i & 174 + 76i \\ 4329 - 262i & -643 + 39i & 12753 - 771i & -58 + 6i \end{pmatrix}$$

The eigenvalues are  $-2i$ ,  $-i$ ,  $3i/2$  and  $3/2$ .

Let us fix the accuracy at  $\varepsilon = 10^{-6}$ . The last 7 iterations of  $\lambda_1^{(k)}$  are:

0.0000477588150960872 - 1.99991424541241 *i*  
 -0.0000479821875446196 - 1.99998019901599 *i*  
 -0.0000272650944159076 - 2.00002375338328 *i*  
 0.0000271520045767515 - 2.00002973125038 *i*  
 0.0000154506695115737 - 1.99997272532314 *i*  
 -0.0000152424622193764 - 1.99999349337182 *i*

Power method - demonstration in  $\mathbb{C}^{n,n}$  (2/2)

# Power method: other eigenvalues

Suppose that using the power method we found the dominant eigenvalue  $\lambda_1$  and its corresponding (normalized) eigenvector  $u_1$ . How can we find the other eigenvalues?

# Power method: other eigenvalues

Suppose that using the power method we found the dominant eigenvalue  $\lambda_1$  and its corresponding (normalized) eigenvector  $u_1$ . How can we find the other eigenvalues?

Suppose that the matrix  $M$  is **normal** (i.e., that  $MM^* = M^*M$ , where  $M^*$  is the conjugate transpose of  $M$ ). Then its eigenvectors are orthogonal.

# Power method: other eigenvalues

Suppose that using the power method we found the dominant eigenvalue  $\lambda_1$  and its corresponding (normalized) eigenvector  $u_1$ . How can we find the other eigenvalues?

Suppose that the matrix  $M$  is normal (i.e., that  $MM^* = M^*M$ , where  $M^*$  is the conjugate transpose of  $M$ ). Then its eigenvectors are orthogonal.

We can consider a new matrix  $M'$  defined as:

$$M' := M - \lambda_1 u_1 \cdot u_1^*$$

# Power method: other eigenvalues

Suppose that using the power method we found the dominant eigenvalue  $\lambda_1$  and its corresponding (normalized) eigenvector  $u_1$ . How can we find the other eigenvalues?

Suppose that the matrix  $M$  is **normal** (i.e., that  $MM^* = M^*M$ , where  $M^*$  is the conjugate transpose of  $M$ ). Then its eigenvectors are orthogonal.

We can consider a new matrix  $M'$  defined as:

$$M' := M - \lambda_1 u_1 \cdot u_1^*$$

The matrix  $M'$  has  $u_1$  as eigenvector, but the associated eigenvalue is 0, indeed:

$$M' u_1 = M u_1 - \lambda_1 u_1 \cdot \|u_1\|^2 = \lambda_1 u_1 - \lambda_1 u_1 = 0.$$

# Power method: other eigenvalues

Suppose that using the power method we found the dominant eigenvalue  $\lambda_1$  and its corresponding (normalized) eigenvector  $u_1$ . How can we find the other eigenvalues?

Suppose that the matrix  $M$  is **normal** (i.e., that  $MM^* = M^*M$ , where  $M^*$  is the conjugate transpose of  $M$ ). Then its eigenvectors are orthogonal.

We can consider a new matrix  $M'$  defined as:

$$M' := M - \lambda_1 u_1 \cdot u_1^*$$

The matrix  $M'$  has  $u_1$  as eigenvector, but the associated eigenvalue is 0, indeed:

$$M' u_1 = M u_1 - \lambda_1 u_1 \cdot \|u_1\|^2 = \lambda_1 u_1 - \lambda_1 u_1 = 0.$$

We can now apply the power method to the matrix  $M'$ . The dominant eigenvalue of  $M'$  will be the second largest (in absolute value) eigenvalue of  $M$ .

# Outline

- 1 Eigenvectors
- 2 Power method
- 3 QR algoritmus



# QR factorization and QR algorithm (1/2)

The power method is not suitable to find all eigenvalues of a given matrix  $M$ .

# QR factorization and QR algorithm (1/2)

The power method is not suitable to find all eigenvalues of a given matrix  $M$ .

Other algorithms are based on the fact that similar matrices have the same eigenvalues. The goal of QR algorithm is to construct a sequence  $(M_k)_{k=0}^{\infty}$  of similar matrices in the following way:

$$M_0 = M \quad \text{and} \quad M_k = P_k M_{k-1} P_k^{-1} \quad k \in \mathbb{N},$$

where each  $P_k$  is a regular matrix,  $M_k \rightarrow M_{\infty}$  and for  $M_{\infty}$  is easy to find the eigenvalues (for instance,  $M_{\infty}$  is upper triangular).

## QR factorization and QR algorithm (2/2)

The **QR factorization** consists in expressing a real (or complex) matrix  $M \in \mathbb{R}^{n,n}$  as a product

$$M = Q \cdot R$$

where  $Q$  is an orthogonal matrix (unitary if  $M \in \mathbb{C}^{n,n}$ ) and  $R$  is upper triangular.

## QR factorization and QR algorithm (2/2)

The **QR factorization** consists in expressing a real (or complex) matrix  $M \in \mathbb{R}^{n,n}$  as a product

$$M = Q \cdot R$$

where  $Q$  is an orthogonal matrix (unitary if  $M \in \mathbb{C}^{n,n}$ ) and  $R$  is upper triangular.

There exist several algorithms to compute such a factorization (Gram-Schmidt, LR algorithm, ...)

## QR factorization and QR algorithm (2/2)

The **QR factorization** consists in expressing a real (or complex) matrix  $M \in \mathbb{R}^{n,n}$  as a product

$$M = Q \cdot R$$

where  $Q$  is an orthogonal matrix (unitary if  $M \in \mathbb{C}^{n,n}$ ) and  $R$  is upper triangular.

There exist several algorithms to compute such a factorization (Gram-Schmidt, LR algorithm, ...)

The **QR algorithm** consists in applying such a factorization at any step, that is for every  $k$  we have

$$M_k = Q_k \cdot R_k$$

and we define

$$M_{k+1} := R_k Q_k = Q_k^{-1} Q_k R_k Q_k = Q_k^{-1} M_k Q_k.$$

We start the iteration with  $M_0 = M$ . Every matrix  $M_k$  is similar to the previous matrix  $M_{k-1}$  in the sequence, so that all matrices have the same eigenvalues. Under certain conditions  $M_k$  converges to a triangular matrix.