

Part I

Quick outline of the course

Organization

 Organization


Lecturer:

1. Francesco Dolce (dolcefra@fit.cvut.cz)

 Conditions, materials, schedules: <https://courses.fit.cvut.cz/NIE-MPI/>

 see [here](#) the conditions to pass the course

Why mathematics?

 Why mathematics?


 Why should we learn mathematics?



If someone can take up this position (painlessly),
what do you say to yourself? [0.2cm] *Good! I'd like to be agile as she is ...*
[0.2cm] OR [0.2cm] *Hmm, I didn't need such a daredevil position in my life,
I am going to train sitting on a chair instead, that's what I do ...*

Understanding

MATHEMATICS
is not about
numbers, equations,
computations, or
algorithms:
it is about
UNDERSTANDING.

William Paul Thurston

15 Majors that
Will Make You
Rich (measured
by money)

1. Petroleum Engineering (\$155,000 – after some time)
2. Physics (\$101,800)
3. Applied Mathematics (\$98,600 “Jobs in this field can be found in nearly every sector.”)
4. Computer Science (\$97,900)
5. Biomedical Engineering (\$97,800)
6. Statistics (\$93,800)
7. Civil Engineering (\$90,200)
8. Mathematics (\$89,900)
9. Environmental Engineering (\$88,600)
10. Software Engineering (\$87,800)
11. Finance (\$87,300)
12. Construction Management (\$85,200)
13. Biochemistry (\$84,700)
14. Geology (\$83,300)
15. Management Information Systems (\$82,200)

source: <http://likes.com/misc/15-majors-that-will-make-you-rich>

Famous names

George STIBITZ (Ph.D. in mathematical physics)


He was a Bell Labs researcher known for his work in the 1930s and 1940s on the realization of Boolean logic digital circuits using electromechanical relays as the switching element.



 Famous names
 ...

Marian REJEWSKI, Alan TURING, ... (mathematicians)

Breaking of German codes during WWII.



 Famous names
 ...

Claude SHANNON, (founder of information theory, mathematician)

Shannon is famous for having founded information theory with one landmark paper published in 1948. But he is also credited with founding both digital computer and digital circuit design theory in 1937, when, as a 21-year-old master's student at MIT, he wrote a thesis demonstrating that electrical application of Boolean algebra could construct and resolve any logical, numerical relationship.



Famous names

...

Dennis RITCHIE, (computer scientist, creator of C programming language)

Ritchie graduated from Harvard University with degrees in physics and applied mathematics.



Famous names

...

Linus TORVALDS (developer of the Linux kernel)


His parents were both journalists. However, he was highly influenced by his maternal grandfather to pursue his career in computers. Since childhood, Linus **was brilliant in mathematics**. In 1988 he began studying **computer science** at the University of Helsinki. Linus is from a minority group in Finland and his first language is not Finnish but Swedish. For this reason, his pronunciation of Linux in Swedish were not understood or often taken as an error.



 Famous names
 ...

Bill GATES (founder of Microsoft)

In his sophomore year, Gates **devised an algorithm for pancake sorting** as a solution to one of a series of unsolved problems presented in a combinatorics class by Harry Lewis, one of his professors. Gates' solution held the record as the fastest version for over thirty years; its successor is faster by only 2%. His solution was later formalized in a published paper in collaboration with Harvard computer scientist Christos Papadimitriou.




 Famous names
 ...

Larry PAGE and Sergey BRIN (founders of Google)

Larry was in search of a dissertation theme for his PhD in computer science and considered exploring the **mathematical properties of the World Wide Web**, understanding its link structure as a huge graph.

After graduation at the University of Maryland, Sergey moved to Stanford University to acquire a Ph.D in computer science.

The company was founded while they were both attending Stanford University.




Short overview of included topics

What about us?

What will we be talking about in this course?

Multivariate
functions and
optimization

- Many problems can be formulated as optimization problems: we maximize/minimize some functions that determines gain/cost/time/distance ...
- If the function is given analytically, we know how to find the optimum.



$$[2\text{mm}] \sin(x \cdot y)$$

General algebra

Notions from general algebra are one of the basic mathematical tools.

·	1	2	3	4	5	6	7	8	9	10	11	12
1	1	2	3	4	5	6	7	8	9	10	11	12
2	2	4	6	8	10	12	1	3	5	7	9	11
3	3	6	9	12	2	5	8	11	1	4	7	10
4	4	8	12	3	7	11	2	6	10	1	5	9
5	5	10	2	7	12	4	9	1	6	11	3	8
6	6	12	5	11	4	10	3	9	2	8	1	7
7	7	1	8	2	9	3	10	4	11	5	12	6
8	8	3	11	6	1	9	4	12	7	2	10	5
9	9	5	1	10	6	2	11	7	3	12	8	4
10	10	7	4	1	11	8	5	2	12	9	6	3
11	11	9	7	5	3	1	12	10	8	6	4	2
12	12	11	10	9	8	7	6	5	4	3	2	1


[1mm] Cayley

table of the group \mathbb{Z}_{13}^\times


Besides a general introduction, we will focus on finite groups and fields, which form the basis for cryptography, hash functions, etc.

Fuzzy Logic

Describe systems by properties which are not evaluated by values beyond just **true** or **false**.

Numerical
mathematics

Continuous mathematics using the computer, stability of numerical algorithms ...



Part II

Multivariate optimisation

Examples

Problem 1. *Imagine the following situation: You have created a program that processes a text input by a user. You know, from theoretical analysis of the source code and algorithms used within the program, that it is impossible to determine the exact time needed to process a text of length k . However, you know that it is **approximately** proportional to the length of the text.*

Duration of a text processing program (1 of 6)

Mathematically: Denote $t(k)$ the “average” number of seconds needed to process a text of length k . We know that

$$t(k) \approx \alpha k \quad \text{for some } \alpha \in \mathbb{R}.$$

Problem: The **proportionality constant** α is unknown. How would you reasonably estimate its value?

Sketch of a solution:

Duration of a text processing program (2 of 6)

1. Run the program for several, say n , texts of various lengths and measure the actual running times. This gives us n couples of measurements $(k_1, t_1), (k_2, t_2), \dots, (k_n, t_n)$.
2. For a given α , we can measure the approximation error $t(k) \approx \alpha k$ by this function:

$$e(\alpha) = (t_1 - \alpha k_1)^2 + (t_2 - \alpha k_2)^2 + \dots + (t_n - \alpha k_n)^2 = \sum_{i=1}^n (t_i - \alpha k_i)^2.$$

3. In order to find the best approximating proportionality constant α , we find the value of α for which the error $e(\alpha)$ is minimal:

an optimal value of α is a minimum point of the function $e(\alpha)$.

How to find a minimum point of $e(\alpha)$:

Duration of a text processing program (3 of 6)

1. Find the **first derivative** $e'(\alpha)$:

$$e'(\alpha) = \left(\sum_{i=1}^n (t_i - \alpha k_i)^2 \right)' = \sum_{i=1}^n -2k_i(t_i - \alpha k_i).$$

2. Find the **critical points**, i.e., the points α_0 where $e'(\alpha_0)$ is zero or does not exist:

$$e'(\alpha_0) = 0 \Leftrightarrow \sum_{i=1}^n -2k_i(t_i - \alpha_0 k_i) = 0 \Leftrightarrow \sum_{i=1}^n k_i t_i = \alpha_0 \sum_{i=1}^n k_i^2 \Leftrightarrow \alpha_0 = \frac{\sum_{i=1}^n k_i t_i}{\sum_{i=1}^n k_i^2}$$

3. The critical points are our candidates for the points of (local) minimal or maximal values of the function e . To be sure that the value of α we found is a minimum we need the **second derivative**:

$$e''(\alpha) = \left(\sum_{i=1}^n -2k_i(t_i - \alpha k_i) \right)' = \sum_{i=1}^n 2k_i^2.$$

...continues ...

Duration of a
text processing
program (4 of 6)

We know that if $e''(\alpha_0) > 0$ (resp. $e''(\alpha_0) < 0$), then the critical point α_0 is a local strict minimum (resp. strict maximum) point.

If $e''(\alpha_0) = 0$, then α is neither of these two cases (maybe an inflexion point?).

Solution: based on our measurements $(k_1, t_1), (k_2, t_2), \dots, (k_n, t_n)$, we get the best approximation $t(k) \approx \alpha k$ for

$$\alpha = \alpha_0 = \frac{\sum_{i=1}^n k_i t_i}{\sum_{i=1}^n k_i^2}.$$

Indeed, this α_0 is the unique (why unique?) global (why global?) minimum point of the approximation error function $e(\alpha)$ since the second derivative

$$e''(\alpha_0) = \sum_{i=1}^n 2k_i^2 \text{ is positive.}$$

Duration of a
text processing
program (5 of 6)

Problem 2 (slight modification). *Imagine the following situation: You have created a program that processes a text input by a user. You know, from theoretical analysis of the source code and algorithms used within the program, that it is impossible to determine precisely the time needed to process a text of length k . However, you know that it is **approximately** proportional to the length of the text **and to the frequency of the processor**.*

Mathematically: Denote by $t(k, f)$ the “average” number of seconds needed to process a text of length k , and the frequency of the processor by f . We know that

$$t(k, f) \approx \alpha k + \beta f \quad \text{for some } \alpha, \beta \in \mathbb{R}.$$

Problem: The constants α and β are unknown. How would you reasonably estimate their values?

Duration of a
text processing
program (6 of 6)

Sketch of solution:

1. Run the program for several, say n , texts of various lengths on computers with different frequencies and measure the actual running times. This gives us n triplets of measurements $(k_1, t_1, f_1), (k_2, t_2, f_2), \dots, (k_n, t_n, f_n)$.
2. For a given α and β , we can measure the approximation error $t(k, f) \approx \alpha k + \beta f$ by this **two-variable** function:

$$\begin{aligned} e(\alpha, \beta) &= (t_1 - \alpha k_1 - \beta f_1)^2 + (t_2 - \alpha k_2 - \beta f_2)^2 + \dots + (t_n - \alpha k_n - \beta f_n)^2 = \\ &= \sum_{i=1}^n (t_i - \alpha k_i - \beta f_i)^2. \end{aligned}$$

3. In order to find the best approximating constants α and β , we find values of α and β for which the error $e(\alpha, \beta)$ is minimal: an optimal value of α and β is the “two-dimensional” minimum point of $e(\alpha, \beta)$.

Comments

Why “optimization”?

A typical situation in physics, engineering, economy, chemistry, etc. is that you have a function that measures your profit, your loss, the energy of something, etc.

The value of such function is given by one or more inputs and the relation between inputs and the resulting value is usually stated as a mathematical formula since all these sciences uses mathematical models to understand and quantify their subject of interest.


An example of such function is our function $e(\alpha, \beta)$ that measures the approximation error.

Typically, we want to maximize or minimize such functions (maximize the profit, the energy, minimize the loss, the error) which leads to the problem of finding **optimal** values of the inputs. Therefore the name “optimization”.

Comments

There is **another very important usage** of the derivative.

Derivatives measure the rate of change of a function. This helps us to describe the behaviour of a **dynamical systems** like a ball on a spring:



The position of the ball at time t is a function $x(t)$ satisfying the differential equation

$$x''(t) + \omega^2 x(t) = 0.$$

The solution of this equation is

$$x(t) = x_0 \cos(\omega t) + \frac{v_0}{\omega} \sin(\omega t), \quad t \in \mathbb{R},$$

where $x_0 = x(0)$ and v_0 are the position and the speed of the ball at time $t = 0$. This model is known as **harmonic oscillator**.

Univariate optimization

Derivative

How do we differentiate?

Example 3. Find the first derivative of $f(x)$, where

(a) $f(x) = x^3 + 4x^2 + 6$,

(b) $f(x) = \sin(x^3)$,


(c) $f(x) = e^x \sin x$.

Solutions:

(a) $f'(x) = 3x^2 + 8x$,

(b) $f'(x) = 3x^2 \cos(x^3)$,

(c) $f'(x) = e^x \sin x + e^x \cos x$.



Geometrical
meaning of
derivative:
tangent line (1
of 2)

- The tangent line to the graph of a function $f(x)$ at a point x_0 is a straight line that “just touches” the curve at that point.
- Any straight line has equation $y = ax + b$, where a is the **slope** of the line.
- **The slope of the tangent line to $f(x)$ at the point x_0 equals the first derivative evaluated at x_0 : $f'(x_0)$.**
- The tangent line at the point x_0 satisfies the equation

$$y = f'(x_0)(x - x_0) + f(x_0).$$

Geometrical
meaning of
derivative:
tangent line (2
of 2)

With this geometrical explanation it is easy to see that the following statements are true:

- If $f'(x_0)$ is positive, then $f(x)$ is increasing at x_0 .
- If $f'(x_0)$ is negative, then $f(x)$ is decreasing at x_0 .
- If x_0 is a local minimum/maximum point of $f(x)$, then $f'(x_0) = 0$ or $f'(x_0)$ does not exist. Such points are called **critical points**.

Example 4. Find all critical points of

$$f(x) = \frac{x^3}{3} + 2x^2 + 3x + 6.$$

What does it mean that the second derivative $f''(x)$ is positive?

- The second derivative is a derivative of the first derivative; therefore the fact that $f''(x)$ is positive implies that $f'(x)$ is increasing (at the point x).
- If $f'(x)$ is increasing, then the function $f(x)$ is more and more increasing (if $f'(x) > 0$) or less and less decreasing (if $f'(x) < 0$).
- An illustrative example of function with positive second derivative is $f(x) = x^2$.

Again, if we understand the geometrical meaning of the second derivative, we can easily see that the following statements are true:

Theorem 5. Let x_0 be a critical point of a function $f(x)$ such that $f'(x_0) = 0$ and $f''(x_0)$ exists.

- If $f''(x_0) > 0$, then the function is **convex** at x_0 and x_0 is a point of a (strict) minimum.
- If $f''(x_0) < 0$, then the function is **concave** at x_0 and x_0 is a point of a (strict) maximum.

Question: what can happen if $f''(x_0) = 0$?

Given a function $f(x)$, we want to find its extremal values.


1. Find the first derivative $f'(x)$.
2. Find the critical points: solve the equation $f'(x) = 0$ and find the points where the derivative **does not exist**.
3. Find the second derivative $f''(x)$.
4. If possible, for all critical points x_0 evaluate $f''(x_0)$ and decide whether this point is a point of minimum or maximum or whether it is an inflexion point. (Other critical points have to be treated by hand.)

The goal of this and the next lecture is to understand what happens when we have more than 1 variable. We shall build a similar cookbook for such functions.

Multivariate optimization

For a univariate function $f(x)$, its graph is the set of points $(x, f(x))$ which can be depicted in Cartesian coordinate system (typically with x - and y -axis).
[2mm]


What if the function depends on more variables? For instance: $f(x, y)$.



Graph of a two-variable function $\sin(x \cdot y)$: the set of points $(x, y, \sin(x \cdot y))$.

- To depict a graph of a two-variable function we need a third axis (typically z -axis) and a 3-dimensional figure. Such graph is in general some surface.
- It is impossible to (directly) depict graphs of functions of more than 2 variables since we cannot make 4 or more dimensional figures.

Example 6. How does the graph of $f(x, y) = x^2 - y^2$ look like?



Given the function $f(x, y) = x^2 + xy + y^2$.

- If we fix the value of the variable y to 3, we obtain a univariate function $f(x) = x^2 + 3x + 9$ having its derivative equal to $2x + 3$.
- We can fix the value of y not only to a specific number: **we just treat y as a constant**. Then we get univariate function $f(x) = x^2 + xy + y^2$ and its derivative is $2x + y$.
- This derivative is called **partial derivative of (x, y) with respect to x** and denoted by

$$\frac{\partial f}{\partial x}(x, y) = 2x + y.$$

- In the same way we define the partial derivative of $f(x, y)$ with respect to y :

$$\frac{\partial f}{\partial y}(x, y) = x + 2y.$$

- In general $\frac{\partial f}{\partial x}(x, y)$ and $\frac{\partial f}{\partial y}(x, y)$ are two-variate functions.

The derivative of a (single variate) function $f(x)$ is the following limit (if it exists):

$$f'(x) = \lim_{\delta \rightarrow 0} \frac{f(x + \delta) - f(x)}{\delta}.$$

Partial derivatives are defined similarly:

Definition 7. *The partial derivative of $f(x_1, x_2, \dots, x_n)$ with respect to x_i at the point (x_1, x_2, \dots, x_n) is defined by (if the limit below exists)*

$$\begin{aligned} \frac{\partial f}{\partial x_i}(x_1, x_2, \dots, x_{i-1}, x_i, x_{i+1}, \dots, x_n) &= \\ &= \lim_{\delta \rightarrow 0} \frac{f(x_1, x_2, \dots, x_{i-1}, x_i + \delta, x_{i+1}, \dots, x_n) - f(x_1, x_2, \dots, x_n)}{\delta}. \end{aligned}$$


Since the definition is similar, even the geometrical meaning is analogous.

The partial derivatives of $f(x, y)$ can be in short denoted by

$$f_x(x, y) = \frac{\partial f}{\partial x}(x, y) \quad \text{and} \quad f_y(x, y) = \frac{\partial f}{\partial y}(x, y).$$

The number $f_x(x, y)$ for given values of x and y is again the slope of a tangent line, but a surface has infinitely many tangent lines in all possible directions at any point, so which one is this one?

It is the only tangent line which is parallel to the x -axis.



Definition 8. For a function $f(x_1, x_2, \dots, x_n)$ we define second partial derivatives

$$f_{x_j x_i}(x_1, x_2, \dots, x_n) = \frac{\partial^2 f}{\partial x_j \partial x_i}(x_1, x_2, \dots, x_n) = \frac{\partial}{\partial x_j} \left(\frac{\partial f}{\partial x_i}(x_1, x_2, \dots, x_n) \right),$$

in particular, for $i = j$ we have

$$f_{x_i x_i}(x_1, x_2, \dots, x_n) = \frac{\partial^2 f}{\partial x_i^2}(x_1, x_2, \dots, x_n) = \frac{\partial}{\partial x_i} \left(\frac{\partial f}{\partial x_i}(x_1, x_2, \dots, x_n) \right).$$

Example 9. Find partial derivatives with respect to all variables

(a) $f(x, y) = xy + e^x \cos y,$

(b) $f(x, y) = x^2 y^3 + x^3 y^4 - e^{xy^2},$

(c) $f(x, y, z) = \sin(xy/z).$

Example 10. Find all second partial derivatives of the functions

(a) $f(x, y) = x^2 + xy^2 + 3x^3 y,$

(b) $f(x, y, z) = e^{xz} + y \cos x,$

(c) $f(x, y, z) = z \cos(xy) + x \sin(yz).$

The fact that the mixed partial derivatives are equal is not a coincidence:

Theorem 11. If a function $f(x, y)$ has continuous second partial derivatives, then the mixed second derivatives are equal, i.e.,

$$\frac{\partial^2 f}{\partial y \partial x} = \frac{\partial^2 f}{\partial x \partial y}.$$

This theorem is not true in general, a counterexample is the function

$$f(x, y) = \begin{cases} 0 & \text{at point } (0, 0) \\ \frac{xy(x^2 - y^2)}{x^2 + y^2} & \text{otherwise.} \end{cases}$$

