

# NIE-MPI: Tutorial 9

created: September 11, 2024, 15:09

## 9.1 Numerical mathematics

### Machine numbers

If not stated otherwise, we consider **single precision** and mathematical rounding.

**Exercise 9.1.** Are the following numbers machine numbers?

- a)  $10^{113}$
- b)  $1 + 2^{-32}$
- c)  $\frac{1}{5}$
- d)  $\frac{3}{10}$
- e)  $\frac{3}{256}$
- f)  $2^{-20} - 16$

**Exercise 9.2.** Which number correspond, in base 10, to the following machine numbers? (we use the notation (**sign, mantissa, exponent**))

- a) (1, 101000000000000000000000, 01000010);
- b) (0, 101000000000000000000000, 00000001);
- c) (1, 101000000000000000000000, 00000000).

**Exercise 9.3.** Which are the closest neighbours, between the normalized machine numbers, of the normalized machine number  $2^t$ ?

**Exercise 9.4.** Consider a decimal machine in which 2 decimal digits are allocated to the significand (and we do not care about the exponent) and the approximation is done by rounding. Sum the following numbers 0,25, 0,0034, 0,00051 a 0,061 in the following order:

- a) from the least to the greatest,
- b) from the greatest to the least.

Compare to the exact result.

**Exercise 9.5.** Find the absolute and the relative errors of the following pair  $(a, \alpha)$ , where  $\alpha$  is an approximate value of  $a$ :

- $(0.100, 0.110)$ ,
- $(0.100, 0.101)$ ,
- $(0, 500, 0.510)$ ,
- $(0.500, 0.501)$ .

**Exercise 9.6.** Let  $x$  and  $y$  be normalized machine numbers. Which following statements are true if we suppose that no underflow or overflow happens (and we stay within normalized numbers)?

1.  $\text{fl}(x + y) = \text{fl}(x) + \text{fl}(y)$ ;
2.  $\text{fl}(x + y) = \text{fl}(y + x)$ ;
3.  $\text{fl}((x + y) + z) = \text{fl}(x + (y + z))$ ;

If a statement is not true, find a counterexample. If it is true, give an argument. If your answer depends on something else, mention it.