

MPI - Lecture 6

What shall we do today?

- Multivariate optimization:
 - Gradient
 - Tangent plane
 - Critical points on two or more variables
 - Hessian (matrix)

Gradient


Gradient of a function

The **gradient** of a function $f(x_1, x_2, \dots, x_n)$ at the (n -dimensional) point $b \in \mathbb{R}^n$ is the n -dimensional vector function $\nabla f(b)$ defined by

$$\nabla f(b) = \left(\frac{\partial f}{\partial x_1}(b), \frac{\partial f}{\partial x_2}(b), \dots, \frac{\partial f}{\partial x_n}(b) \right).$$

Example 1. Find the gradient of the function $f(x, y) = x^2 + xy + y^2$ at the point $(1, 1)$.

Geometrical meaning: the gradient points in the direction of the greatest rate of increase of the function. Its magnitude equals the rate of increase.



Gradient and
the directional
derivative

We saw that the partial derivative with respect to x at the point a is equal to the slope of tangent line at this point in direction parallel to the x -axis.

Example 2. If we are on the graph of the function $f(x, y) = x^2 + xy + y^2$ at the point $(1, 1)$ and we start moving in the direction parallel to the x -axis, i.e., in the direction of the vector $(1, 0)$, we will go “uphill” under the angle $\arctan 3$ since

$$\frac{\partial f}{\partial x}(1, 1) = 2 + 1 = 3.$$

What will be the slope if we move in the direction of a general vector $\vec{v}$?


Theorem 3. Given a function $f(x) : \mathbb{R}^n \rightarrow \mathbb{R}$, a point $a \in \mathbb{R}^n$ and a **unit** vector $\vec{v} \in \mathbb{R}^n$, the **derivative in the direction of the vector $\vec{v}$** is the dot product of the gradient and $\vec{v}$, i.e., $\nabla f(a_1, a_2, \dots, a_n) \cdot \vec{v}$.

Tangent plane

Tangent plane

The **tangent plane** to a function $f(x, y)$ at the point (x_0, y_0) is a 2-dimensional plane that “touches” the graph of the function at (x_0, y_0) . Its equation is

$$z = \frac{\partial f}{\partial x}(x_0, y_0) \cdot (x - x_0) + \frac{\partial f}{\partial y}(x_0, y_0) \cdot (y - y_0) + f(x_0, y_0).$$



Example 4. Find the tangent plane to $f(x, y) = x^2 + xy + y^2$ at $(1, 1)$.

Critical points

Critical points –
two variables

- In the one dimensional case the critical points are those points where the tangent line is parallel to the x -axis, i.e., points where $f'(x) = 0$, or where the derivative does not exist.
- The **critical points** of a two variable function are those points where the tangent plane is parallel to the plane given by the x -axis and the y -axis or where the gradient does not exist.

The first class of these points can be found as a solution of

$$\nabla f(x, y) = (0, 0)$$

which leads to the system of two equations for two variables

$$\begin{cases} \frac{\partial f}{\partial x}(x, y) = 0 \\ \frac{\partial f}{\partial y}(x, y) = 0 \end{cases}.$$

Critical points –
more variables

For an n -variable function $f(x_1, x_2, \dots, x_n)$ the situation is analogous:

The **critical points** of $f(x_1, x_2, \dots, x_n)$ are points satisfying the equation

$$\nabla f(x_1, x_2, \dots, x_n) = 0$$

i.e., points satisfying the system of n equations for n variables

$$\begin{cases} \frac{\partial f}{\partial x_1}(x_1, x_2, \dots, x_n) = 0 \\ \frac{\partial f}{\partial x_2}(x_1, x_2, \dots, x_n) = 0 \\ \vdots \\ \frac{\partial f}{\partial x_n}(x_1, x_2, \dots, x_n) = 0 \end{cases},$$

or where the gradient does not exist.

(Instead of a tangent plane, we have a **tangent hyperplane**.)

Critical points –
example

Example 5. Find all critical points of

$$f(x_1, x_2, x_3) = x_1x_3 + x_1^2 - x_2 + x_2x_3 + x_2^2 + 3x_3^2,$$

We get

$$\nabla f(x_1, x_2, x_3) = (x_3 + 2x_1, -1 + x_3 + 2x_2, x_1 + x_2 + 6x_3)$$

which always exists. Thus the critical points are the solution of the system

$$\begin{cases} x_3 + 2x_1 &= 0 \\ -1 + x_3 + 2x_2 &= 0 \\ x_1 + x_2 + 6x_3 &= 0 \end{cases},$$

which, using the standard procedure for a system of linear equations, gives us the only solution $\left(\frac{1}{20}, \frac{11}{20}, \frac{-1}{10}\right)$.

Hessian

Type of a critical point (1 of 4)

In the one dimensional case, we can use the second derivative to decide the type of the critical point.

Theorem 6. Let x_0 be a critical point of a function $f(x)$ such that $f'(x_0) = 0$ and $f''(x_0)$ exists.

- If $f''(x_0) > 0$, then the function is **convex** at x_0 , and x_0 is a point of a minimum.
- If $f''(x_0) < 0$, then the function is **concave** at x_0 , and x_0 is a point of a maximum.
- If $f''(x_0) = 0$, then x_0 may be a minimum, maximum, inflection point, ...

Do we have something similar for more variables? What is the second derivative?

Type of a critical point (2 of 4)

The analogue of the second derivative is the following.

Definition 7. For a function $f(x_1, x_2, \dots, x_n)$ we define the *Hessian matrix* as

$$\nabla^2 f(x_1, x_2, \dots, x_n) = \begin{pmatrix} \frac{\partial^2 f}{\partial x_1^2}(x_1, \dots, x_n) & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n}(x_1, \dots, x_n) \\ \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1}(x_1, \dots, x_n) & \cdots & \frac{\partial^2 f}{\partial x_n^2}(x_1, \dots, x_n) \end{pmatrix}$$

assuming that all the derivatives exist.

Type of a critical point (3 of 4)

We would like to construct rules like “If $f''(x_0) > 0$, then the critical point x_0 is the point of minimum”.

But to say that the matrix is “positive” is problematic ... Let us use a different notion.

Definition 8. A matrix $A \in \mathbb{R}^{n,n}$ is

- (i) *positively definite* if for all non-zero vectors $a \in \mathbb{R}^n$ it holds that $aAa^T > 0$;
- (ii) *positively semidefinite* if for all vectors $a \in \mathbb{R}^n$ it holds that $aAa^T \geq 0$ and the equality is true for at least one non-zero vector $b \in \mathbb{R}^n$;
- (iii) *negatively definite* if for all non-zero vectors $a \in \mathbb{R}^n$ it holds that $aAa^T < 0$;
- (iv) *negatively semidefinite* if for all vectors $a \in \mathbb{R}^n$ it holds that $aAa^T \leq 0$ and the equality is true for at least one non-zero vector $b \in \mathbb{R}^n$;
- (v) *indefinite* otherwise.

Type of a critical point (4 of 4)

Theorem 9. If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ has all second partial derivative continuous at a critical point $b \in \mathbb{R}^n$, then

- (i) if $\nabla^2 f(b)$ is positively definite, then b is a point of local minimum;
- (ii) if $\nabla^2 f(b)$ is negatively definite, then b is a point of local maximum;

(iii) if $\nabla^2 f(b)$ is indefinite, then b is a saddle point.

Sylvester's criterion on definiteness

For an $n \times n$ dimensional **symmetric** matrix A we define the **principal minors**:

- M_1 is the upper left 1-by-1 corner of A ,
- M_2 is the upper left 2-by-2 corner of A ,
- ...
- M_n is the upper left n -by- n corner of A .

Theorem 10. Let $A \in \mathbb{R}^{n,n}$ be a symmetric matrix.

- A is positively definite if and only if the determinants of all principal minors are positive.
- A is negatively definite if and only if the determinant of M_i is negative for odd i and positive for even i .

Example

Example 11. Find all minima and maxima of the function

$$f(x, y) = \frac{3x^4 - 4x^3 - 12x^2 + 18}{12(1 + 4y^2)}.$$

Solution: The critical points are $(-1, 0)$, $(0, 0)$ and $(2, 0)$; they are a saddle point, a point of maximum and a point of minimum, respectively.

