

Interval exchanges, admissibility and branching Rauzy induction

Francesco Dolce¹, Dominique Perrin²

¹ LaCIM, Université du Québec à Montréal, LaCIM
² LIGM, Université Paris-Est

Abstract

We introduce a definition of admissibility for subintervals in interval exchange transformations. We characterize the admissible intervals using a branching version of the Rauzy induction. Using this notion, we prove a property of the natural codings of interval exchange transformations, namely that any derived set of a regular interval exchange set is a regular interval exchange set with the same number of intervals. Derivation is taken here with respect to return words. We also study the case of regular interval exchange transformations defined over a quadratic field and show that the set of factors of such a transformation is primitive morphic. The proof uses an extension of a result of Boshernitzan and Carroll.

Contents

1	Preliminaries	3
1.1	Words and recurrent sets	3
1.2	Return words and derived sets	4
2	Interval exchanges	6
2.1	Interval exchange transformations	6
2.2	Natural coding	9
2.3	Induced transformations and admissible semi-intervals	11
2.4	A closure property	16
3	Rauzy induction	17
3.1	One-sided Rauzy induction	17
3.2	Branching induction	19
3.3	Equivalence relation	21
3.4	Induction and automorphisms	24
4	Interval exchange over a quadratic field	27
4.1	Complexities	28
4.2	Return times	29

4.3	Reduced complexity of admissible semi-intervals	30
4.4	Primitive morphic sets	31

Introduction

Interval exchange transformations were introduced by Oseledec [23] following an earlier idea of Arnold [1]. Interval exchange transformations have been generalized to transformations called linear involutions by Danthony and Nogueira in [13] (for other generalizations, see [25]).

The natural coding of interval exchange produces sequences of linear complexity, including Sturmian sequences, and this has been widely studied (see, for example [15] or [2] for small alphabets).

Rauzy has introduced in [24] a transformation, now called Rauzy induction (or Rauzy-Veech induction), which operates on interval exchange transformations. It actually transforms an interval exchange transformation into another one operating on a smaller interval. Its iteration can be viewed as a generalization of the continued fraction expansion. The induction consists in taking the first return map of the transformation with respect to a subinterval of the interval on which the exchange is defined. The induced map of an interval exchange on s intervals is still an interval exchange with at most $s + 2$ intervals.

Rauzy introduced in [24] the definition of right-admissibility for an interval and characterized the right-admissible intervals as those which can be reached by the Rauzy induction.

Interval exchange transformations defined over quadratic fields have been studied by Boshernitzan and Carroll ([11] and [10]). Under this hypothesis, they showed that, using iteratively the first return map on one of the intervals exchanged by the transformation, one obtains only a finite number of different new transformations up to rescaling, extending the classical Lagrange's theorem that quadratic irrationals have a periodic continued fraction expansion.

In this paper, we generalize both the notion of admissible intervals and of Rauzy induction to a two-sided version.

Our main result is a characterization of the admissible intervals (Theorem 26). We show that, in particular, intervals associated with factors of the natural coding of an interval exchange transformation are admissible (Proposition 20).

Our motivation is the study of the natural coding of these transformations by words, in the spirit of the research initiated in [4] and containing a series of other papers (see for instance [7] and [5] or [6] where interval exchanges are also studied).

We prove a property of the natural codings of regular interval exchange transformations (Theorem 23) saying that the family of these sets of words is closed by derivation, an operation consisting in taking the first return words to a given word as a new alphabet.

We pay special attention to the case of interval exchange transformations defined over a quadratic field. We prove that the family of transformations

obtained from a regular interval exchange transformation by two-sided Rauzy induction is finite up to rescaling. Moreover, we show that the related interval exchange set is obtained as the set of factors of a primitive morphic word.

The paper is organized as follows.

In Section 1 we recall some basic definitions concerning words and sets. Return words and first return words are also introduced.

In Section 2, we give some notions concerning interval exchange transformations. We recall the result of Keane [19] which proves that regularity is a sufficient condition for the minimality of such a transformation (Theorem 7). We also introduce the natural codings of interval exchange transformations. We define the derivate of an interval exchange set with respect to a coding morphism and we show a closure property of these sets.

In Section 3, we first recall the notion of *Rauzy induction* introduced in [24]. We introduce a branching version of Rauzy induction. We prove the generalization of Rauzy's theorems to the two-sided case (Theorems 21 and 26).

In Section 4 we generalize the result of Boshernitzan and Carroll [10], enlarging the family of transformations obtained using induction on every admissible semi-interval. This contains the results of [10] because every semi-interval exchanged by a transformation is admissible, while for $n > 2$ there are admissible semi-intervals that we cannot obtain using the induction only on the exchanged ones. We conclude the Section showing that regular quadratic interval exchange sets are primitive morphic (Theorem 53).

Acknowledgement. This work was supported by grants from Région Île-de-France and ANR project Equinocs.

1 Preliminaries

In this section, we first recall some definitions concerning words. We recall the definition of recurrent and uniformly recurrent sets of words (see [21] for a more detailed presentation). We introduce the notion of first return words and derived words. Derived words have been widely studied, in particular in the context of substitutive dynamics (see [14] for example) and are intimately connected with induction.

1.1 Words and recurrent sets

Let A be a finite nonempty alphabet. All words considered below, unless stated explicitly, are supposed to be on the alphabet A . We denote by A^* the set of all words on A . We denote by ε the empty word and by $A^+ = A^* \setminus \{\varepsilon\}$. We denote by $|w|$ the length of a word w . A set of words is said to be *factorial* if it contains the factors of its elements.

A *morphism* $f : A^* \rightarrow B^*$ is a monoid morphism from A^* into B^* . If $a \in A$ is such that the word $f(a)$ begins with a and if $|f^n(a)|$ tends to infinity with

n , there is a unique infinite word denoted $f^\omega(a)$ which has all words $f^n(a)$ as prefixes. It is called a *fixed point* of the morphism f .

A morphism $f : A^* \rightarrow A^*$ is called *primitive* if there is an integer k such that for all $a, b \in A$, the letter b appears in $f^k(a)$. If f is a primitive morphism, the set of factors of any fixed point of f is uniformly recurrent (see [16] Proposition 1.2.3 for example).

An infinite word y over an alphabet B is called *morphic* if there exists a morphism f on an alphabet A , a fixed point $x = f^\omega(a)$ of f and a morphism $\sigma : A^* \rightarrow B^*$ such that $y = \sigma(x)$. If $A = B$ and σ is the identity map, we call y *purely morphic*. If f is primitive we say that the word is *primitive morphic*.

A factorial set of words $F \neq \{\varepsilon\}$ is *recurrent* if for every $u, w \in F$ there is a $v \in F$ such that $uvw \in F$. For an infinite word x , we denote $F(x)$ the set of factors of x . An infinite word x is *recurrent* if for any $u \in F(x)$ there is a $v \in F(x)$ such that $uvu \in F(x)$. As well known, for any recurrent set F there is a recurrent infinite word x such that $F = F(x)$ and conversely, for any recurrent infinite word x , the set $F(x)$ is recurrent (see for example [20]).

Extending the definition, we say that a set $F(x)$ is *morphic* (resp. *purely morphic*, *primitive morphic*) if the infinite word x is morphic (resp. purely morphic, primitive morphic).

A recurrent set of words F is said to be *uniformly recurrent* if, for any word $u \in F$, there exists an integer $n \geq 1$ such that u is a factor of every word of F of length n .

1.2 Return words and derived sets

Let F be a set of words over an alphabet A . For $w \in F$, let

$$\Gamma_F(w) = \{x \in F \mid wx \in F \cap A^+w\} \quad \text{and} \quad \Gamma'_F(w) = \{x \in F \mid xw \in F \cap wA^+\}$$

be respectively the set of *right return words* and *left return words* to w . When F is recurrent, the sets $\Gamma_S(w)$ and $\Gamma'_S(w)$ are nonempty. Actually both of them are infinite. Let

$$\mathcal{R}_F(w) = \Gamma_F(w) \setminus \Gamma_F(w)A^+ \quad \text{and} \quad \mathcal{R}'_F(w) = \Gamma'_F(w) \setminus A^+\Gamma'_F(w)$$

be respectively the set of *first right return words* and the set of *first left return words* to w . Note that $w\mathcal{R}_F(w) = \mathcal{R}'_F(w)w$.

Clearly, a recurrent set F is uniformly recurrent if and only if the set $\mathcal{R}_F(w)$ (resp. $\mathcal{R}'_F(w)$) is finite for any $w \in F$.

Example 1 Let F be a set of words whose factors of length at most 6 are the labels of the paths starting at the root of the tree represented in Figure 1 (we will see in Example 14 an infinite set of words having such factors).

We have

$$\begin{aligned} \mathcal{R}_F(a) &= \{cbba, ccba, ccbba\}, \\ \mathcal{R}_F(b) &= \{acb, accb, b\}, \\ \mathcal{R}_F(c) &= \{bac, bbac, c\}. \end{aligned}$$

We colored in Figure 1 the words of $\alpha\mathcal{R}_F(\alpha)$ for $\alpha \in A$.




Figure 1: The words of length ≤ 6 of the set F .

For a set of words X and a word u , we denote $u^{-1}X = \{v \in A^* \mid uv \in X\}$. Let F be a recurrent set and let $w \in F$. A *coding morphism* for the set $\mathcal{R}_F(w)$ is a morphism $f : B^* \rightarrow A^*$ which maps bijectively the (possibly infinite) alphabet B onto $\mathcal{R}_F(w)$. The set $f^{-1}(w^{-1}F)$, denoted $\mathcal{D}_f(F)$, is called the *derived set* of F with respect to f . The following result is proved in [7, Proposition 4.3].

Proposition 2 *Let F be a recurrent set. For $w \in F$, let f be a coding morphism for the set $\mathcal{R}_F(w)$. Then*

$$\mathcal{D}_f(F) = f^{-1}(\Gamma_F(w)) \cup \{\varepsilon\}.$$

Let F be a recurrent set and x be an infinite word such that $F = F(x)$. Let $w \in F$ and let f be a coding morphism for the set $\mathcal{R}_F(w)$. Since w appears infinitely often in x , there is a unique factorization $x = vwy$ with $y \in \mathcal{R}_F(w)^\omega$ and v such that vw has no proper prefix ending with w . The infinite word $f^{-1}(y)$ is called the *derived word* of x relative to f , denoted $\mathcal{D}_f(x)$.

Since the set of factors of a recurrent infinite word is recurrent, the following result, proved in [7, Proposition 4.4], shows in particular that the derived set of a recurrent set is recurrent.

Proposition 3 *Let F be a recurrent set and let x be an infinite word such that $F = F(x)$. Let $w \in F$ and let f be a coding morphism for the set $\mathcal{R}_F(w)$. The derived set of F with respect to f is the set of factors of the derived word of x with respect to f , that is $\mathcal{D}_f(F) = F(\mathcal{D}_f(x))$.*

Example 4 Let F be a recurrent set having as factors of length at most 6 the set represented in Figure 1 (we will see such a set in Example 14).

Let f be the coding morphism for the set $\mathcal{R}_F(c)$ given by $f(a) = bac$, $f(b) = bbac$, $f(c) = c$. The derived set of F with respect to f is represented in Figure 2.




Figure 2: The words of length ≤ 3 of $\mathcal{D}_f(F)$.

2 Interval exchanges

In this section we recall the basic definitions of interval exchange transformations, along with minimality and regularity of interval exchanges. We also define the natural coding of an interval exchange and the associated interval exchange set, that is the language of all possible natural codings of a transformation. We then consider the notion of admissibility of a semi-interval. Finally we prove a closure property using the derivation defined in the previous section.

2.1 Interval exchange transformations

Let us recall the definition of an interval exchange transformation (see [12], [28] or [26] for a more detailed presentation).

A *semi-interval* is a nonempty subset of the real line of the form $[\alpha, \beta[= \{z \in \mathbb{R} \mid \alpha \leq z < \beta\}$. Thus it is a left-closed and right-open interval. For two semi-intervals Δ, Γ , we denote $\Delta < \Gamma$ if $x < y$ for any $x \in \Delta$ and $y \in \Gamma$.

Given an order $<$ on A , a partition $(I_a)_{a \in A}$ of a semi-interval $[\ell, r[$ in semi-intervals is *ordered* if $a < b$ implies $I_a < I_b$.

Let now $<_1$ and $<_2$ be two total orders on A . Let $(I_a)_{a \in A}$ be a partition of $[\ell, r[$ in semi-intervals ordered for $<_1$. Let λ_a be the length of I_a . Let $\mu_a = \sum_{b \leq_1 a} \lambda_b$ and $\nu_a = \sum_{b \leq_2 a} \lambda_b$. Set $\alpha_a = \nu_a - \mu_a$. The *interval exchange transformation* relative to $(I_a)_{a \in A}$ is the map $T : [\ell, r[\rightarrow [\ell, r[$ defined by

$$T(z) = z + \alpha_a \quad \text{if } z \in I_a.$$

Observe that the restriction of T to I_a is a translation onto $J_a = T(I_a)$, that μ_a is the right boundary of I_a and that ν_a is the right boundary of J_a . We additionally denote by γ_a the left boundary of I_a and by δ_a the left boundary of J_a . Thus

$$I_a = [\gamma_a, \mu_a[, \quad J_a = [\delta_a, \nu_a[.$$

Since $a <_2 b$ implies $J_a <_2 J_b$, the family $(J_a)_{a \in A}$ is a partition of $[\ell, r[$ ordered for $<_2$. In particular, the transformation T defines a bijection from $[\ell, r[$ onto itself.

An interval exchange transformation relative to $(I_a)_{a \in A}$ is also said to be on the alphabet A . The values $(\alpha_a)_{a \in A}$ are called the *translation values* of the transformation T .

Example 5 Let R be the interval exchange transformation corresponding to $A = \{a, b\}$, $a <_1 b$, $b <_2 a$, $I_a = [0, 1 - \alpha[$, $I_b = [1 - \alpha, 1[$. The transformation R is the rotation of angle α on the semi-interval $[0, 1[$ defined by $R(z) = z + \alpha \bmod 1$.

Since $<_1$ and $<_2$ are total orders, there exists a unique permutation π of A such that $a <_1 b$ if and only if $\pi(a) <_2 \pi(b)$. Conversely, $<_2$ is determined by $<_1$ and π and $<_1$ is determined by $<_2$ and π . The permutation π is said to be *associated* to T .

Set $A = \{a_1, a_2, \dots, a_s\}$ with $a_1 <_1 a_2 <_1 \dots <_1 a_s$. The pair (λ, π) formed by the family $\lambda = (\lambda_a)_{a \in A}$ and the permutation π determines the map T . We will also denote T as $T_{\lambda, \pi}$. The transformation T is also said to be an s -interval exchange transformation.

It is easy to verify that the family of s -interval exchange transformations is closed by taking inverses.

Example 6 Let $T = R^2$ where R is the rotation of Example 5. The transformation T , represented in Figure 3 is a 3-interval exchange transformation. One has $A = \{a, b, c\}$ with $a <_1 b <_1 c$ and $b <_2 c <_2 a$. The associated permutation is the cycle $\pi = (abc)$.




Figure 3: A 3-interval exchange transformation.

The *orbit* of a point $z \in [\ell, r[$ is the set $\mathcal{O}(z) = \{T^n(z) \mid n \in \mathbb{Z}\}$. The transformation T is said to be *minimal* if for any $z \in [\ell, r[$, the orbit of z is dense in $[\ell, r[$.

From now on, set $\gamma_i = \gamma_{a_i}$, $\delta_i = \delta_{a_i}$, $\mu_i = \mu_{a_i}$ and $\nu_i = \nu_{a_i}$. The points $0 = \gamma_1, \mu_1 = \gamma_2, \dots, \mu_{s-1} = \gamma_s$ form the set of *separation points* of T , denoted $\text{Sep}(T)$. Note that the transformation T has at most $s - 1$ *singularities* (that is points at which it is not continuous), which are among the nonzero separation points $\gamma_2, \dots, \gamma_s$.

An interval exchange transformation $T_{\lambda, \pi}$ is called *regular* if the orbits of the nonzero separation points $\gamma_2, \dots, \gamma_s$ are infinite and disjoint. Note that the orbit of 0 cannot be disjoint from the others since one has $T(\gamma_i) = 0$ for some i with $2 \leq i \leq s$.

A regular interval exchange transformation is also said to satisfy the *idoc* condition (where idoc stands for infinite disjoint orbit condition). It is also said to have the Keane property or to be without *connection* (see [9]). As an example,

the 2-interval exchange transformation of Example 5 which is the rotation of angle α is regular if and only if α is irrational.

The following result is due to Keane [19].

Theorem 7 (Keane) *A regular interval exchange transformation is minimal.*

The converse is not true. Indeed, consider the rotation of angle α with α irrational, as a 3-interval exchange transformation with $\lambda = (1 - 2\alpha, \alpha, \alpha)$ and $\pi = (132)$. The transformation is minimal as any rotation of irrational angle but it is not regular since $\mu_1 = 1 - 2\alpha$, $\mu_2 = 1 - \alpha$ and thus $\mu_2 = T(\mu_1)$.

Example 8 Let T be the 3-interval exchange transformation of Example 6 with $\alpha = (3 - \sqrt{5})/2$. The transformation T is regular since α is irrational. Note that $1 - \alpha$ is a separation point which is not a singularity since T is also a 2-interval exchange transformation.

The following necessary condition for minimality of an interval exchange transformation is useful. A permutation π of an ordered set A is called *decomposable* if there exists an element $b \in A$ such that the set B of elements strictly less than b is nonempty and such that $\pi(B) = B$. Otherwise it is called *indecomposable*. If an interval exchange transformation $T = T_{\lambda, \pi}$ is minimal, the permutation π is indecomposable. Indeed, if B is a set as above, the set of orbits of the points in the set $S = \cup_{a \in B} I_a$ is closed and strictly included in $[\ell, r[$. The following example shows that the indecomposability of π is not sufficient for T to be minimal.

Example 9 Let $A = \{a, b, c\}$ and λ be such that $\lambda_a = \lambda_c$. Let π be the transposition (ac) . Then π is indecomposable but $T_{\lambda, \pi}$ is not minimal since it is the identity on I_b .

The iteration of an s -interval exchange transformation is, in general, an interval exchange transformation operating on a larger number of semi-interval.

Proposition 10 *Let T be a regular s -interval exchange transformation. Then, for any $n \geq 1$, T^n is a regular $n(s - 1) + 1$ -interval exchange transformation.*

Proof. Since T is regular, the set $\cup_{i=0}^{n-1} T^{-i}(\mu)$ where μ runs over the set of $s - 1$ nonzero separation points of T has $n(s - 1)$ elements. These points partition the interval $[\ell, r[$ in $n(s - 1) + 1$ semi-intervals on which T is a translation. ■

We close this subsection with a lemma that will be useful in Section 2.

Let T be an interval exchange transformation relative to a partition $(I_i)_{i=1}^s$ and let $(\alpha_i)_{i=1}^s$ be the translations values of T . We say that $\alpha_{m_1} + \alpha_{m_2} + \dots + \alpha_{m_m}$ is an *m -translation value* of T if there exists a point $z_0 \in I_{m_1} \cap T^{-1}(I_{m_2}) \cap \dots \cap T^{-m+1}(I_{m_m})$. Roughly speaking, iterating T we can start from I_{m_1} and arrive to I_{m_m} in exactly m steps, passing (in order) through $I_{m_2}, \dots, I_{m_{m-1}}$.

Moreover, $\alpha_{m_1} + \alpha_{m_2} + \dots + \alpha_{m_m}$ is one of the translation values of the transformation T^m (namely the one corresponding to the semi-interval containing the point z_0).

Note that when T is minimal, every m -translation value of T , with $m > 0$, is different from zero.

Lemma 11 *Let T be a minimal interval exchange transformation over an interval I . For every $N > 0$ there exists an $\varepsilon > 0$ such that for every $z \in I$ and for every $n > 0$, one has*

$$|T^n(z) - z| < \varepsilon \implies n \geq N.$$

Proof. Let $\alpha_1, \alpha_2, \dots, \alpha_s$ be the translation values of T . For every $N > 0$ it is sufficient to choose

$$\varepsilon = \min \left\{ \left| \sum_{j=1}^M \alpha_{i_j} \right| \mid M \leq N \text{ and } \sum_{j=1}^M \alpha_{i_j} \in V_M(T) \right\}.$$

where $V_M(T)$ denotes the set of M -translation values of T . ■

2.2 Natural coding

Let T be an interval exchange transformation relative to $(I_a)_{a \in A}$. For a given real number $z \in [\ell, r[$, the *natural coding* of T relative to z is the infinite word $\Sigma_T(z) = a_0 a_1 \dots$ on the alphabet A defined by

$$a_n = a \quad \text{if} \quad T^n(z) \in I_a.$$

Example 12 Let $\alpha = (3 - \sqrt{5})/2$ and let R be the rotation of angle α on $[0, 1[$ as in Example 5. The natural coding of R relative to α is the Fibonacci word which is the fixed point $t = abaab \dots$ of the morphism f from $\{a, b\}^*$ into itself defined by $f(a) = ab$ and $f(b) = a$ (see, for example, [21, Chapter 2]).

For a word $w = b_0 b_1 \dots b_{m-1}$, let I_w be the set

$$I_w = I_{b_0} \cap T^{-1}(I_{b_1}) \cap \dots \cap T^{-m+1}(I_{b_{m-1}}). \quad (1)$$

Note that each I_w is a semi-interval. Indeed, this is true if w is a letter. Next, assume that I_w is a semi-interval. Then for any $a \in A$, $T(I_{aw}) = T(I_a) \cap I_w$ is a semi-interval since $T(I_a)$ is a semi-interval by definition of an interval exchange transformation. Since $I_{aw} \subset I_a$, $T(I_{aw})$ is a translate of I_{aw} , which is therefore also a semi-interval. This proves the property by induction on the length. The semi-interval I_w is the set of points z such that the natural coding of the transformation relative to z has w as a prefix.

Set $J_w = T^m(I_w)$. Thus

$$J_w = T^m(I_{b_0}) \cap T^{m-1}(I_{b_1}) \cap \dots \cap T(I_{b_{m-1}}). \quad (2)$$

In particular, we have $J_a = T(I_a)$ for $a \in A$. Note that each J_w is a semi-interval. Indeed, this is true if w is a letter. Next, for any $a \in A$, we have

$T^{-1}(J_{wa}) = J_w \cap I_a$. This implies as above that J_{wa} is a semi-interval and proves the property by induction. We set by convention $I_\varepsilon = J_\varepsilon = [0, 1[$. Then one has for any $n \geq 0$

$$a_n a_{n+1} \cdots a_{n+m-1} = w \iff T^m(z) \in I_w \quad (3)$$

and

$$a_{n-m} a_{n-m+1} \cdots a_{n-1} = w \iff T^n(z) \in J_w \quad (4)$$

Let $(\alpha_a)_{a \in A}$ be the translation values of T . Note that for any word w ,

$$J_w = I_w + \alpha_w \quad (5)$$

with $\alpha_w = \sum_{j=0}^{m-1} \alpha_{b_j}$ as one may verify by induction on $|w| = m$. Indeed it is true for $m = 1$. For $m \geq 2$, set $w = ua$ with $a = b_{m-1}$. One has $T^m(I_w) = T^{m-1}(I_w) + \alpha_a$ and $T^{m-1}(I_w) = I_w + \alpha_u$ by the induction hypothesis and the fact that I_w is included in I_u . Thus $J_w = T^m(I_w) = I_w + \alpha_u + \alpha_a = I_w + \alpha_w$. Equation (5) shows in particular that the restriction of $T^{|w|}$ to I_w is a translation.

Note that the semi-interval J_w is the set of points z such that the natural coding of $T^{-|w|}(z)$ has w as a prefix.

If T is minimal, one has $w \in F(\Sigma_T(z))$ if and only if $I_w \neq \emptyset$. Thus the set $F(\Sigma_T(z))$ does not depend on z (as for Sturmian words, see [21]). Since it depends only on T , we denote it by $F(T)$. When T is regular (resp. minimal), such a set is called a *regular interval exchange set* (resp. a minimal interval exchange set).

Let X be the closure of the set of all $\Sigma_T(z)$ for $z \in [\ell, r[$ and let S be the shift on X defined by $S(x) = y$ with $y_n = x_{n+1}$ for $n \geq 0$. The pair (X, S) is a *symbolic dynamical system*, formed of a topological space X and a continuous transformation S . Such a system is said to be minimal if the only closed subsets invariant by S are \emptyset or X . It is well-known that (X, S) is minimal if and only if $F(S)$ is uniformly recurrent (see for example [21] Theorem 1.5.9).

Then we have the following commutative diagram of Figure 4.

$$\begin{array}{ccc} [\ell, r[& \xrightarrow{T} & [\ell, r[\\ \downarrow \Sigma_T & & \downarrow \Sigma_T \\ X & \xrightarrow{S} & X \end{array}$$

Figure 4: A commutative diagram.

The map Σ_T is neither continuous nor surjective. This can be corrected by embedding the interval $[\ell, r[$ into a larger space on which T is a homeomorphism (see [19] or [8] page 349). However, if the transformation T is minimal, the symbolic dynamical system (X, S) is minimal (see [8] page 392). Thus, we obtain the following statement.

Proposition 13 *For any minimal interval exchange transformation T , the set $F(T)$ is uniformly recurrent.*

Note that for a regular interval exchange transformation T , the map Σ_T is injective (see [19] page 30).

Example 14 Let T be the transformation of Example 8. Since T is minimal, the set $F(T)$ is uniformly recurrent. In Subsection 4.4 we will show that the set $F = F(T)$ is the set of factors of the fixed point of a primitive morphism. The words of length at most 6 of the set F are represented in Figure 1.

In Section 4 we will give a sufficient condition for an interval exchange set to be primitive morphic (Theorem 53).

2.3 Induced transformations and admissible semi-intervals

Let T be a minimal interval exchange transformation. Let $I \subset [\ell, r[$ be a semi-interval. Since T is minimal, for each $z \in [\ell, r[$ there is an integer $n > 0$ such that $T^n(z) \in I$.

The *transformation induced* by T on I is the transformation $S : I \rightarrow I$ defined for $z \in I$ by $S(z) = T^n(z)$ with $n = \min\{n > 0 \mid T^n(z) \in I\}$. We also say that S is the *first return map* (of T) on I . The semi-interval I is called the *domain* of S , denoted $D(S)$.

Example 15 Let T be the transformation of Example 8. Let $I = [0, 2\alpha[$. The transformation induced by T on I is

$$S(z) = \begin{cases} T^2(z) & \text{if } 0 \leq z < 1 - 2\alpha \\ T(z) & \text{otherwise.} \end{cases}$$

Let $T = T_{\lambda, \pi}$ be an interval exchange transformation relative to $(I_a)_{a \in A}$. For $\ell < t < r$, the semi-interval $[\ell, t[$ is *right admissible* for T if there is a $k \in \mathbb{Z}$ such that $t = T^k(\gamma_a)$ for some $a \in A$ and

- (i) if $k > 0$, then $t < T^h(\gamma_a)$ for all h such that $0 < h < k$,
- (ii) if $k \leq 0$, then $t < T^h(\gamma_a)$ for all h such that $k < h \leq 0$.

We also say that t itself is right admissible. Note that all semi-intervals $[\ell, \gamma_a[$ with $\ell < \gamma_a$ are right admissible. Similarly, all semi-intervals $[\ell, \delta_a[$ with $\ell < \delta_a$ are right admissible.

Example 16 Let T be the interval exchange transformation of Example 8. The semi-interval $[0, t[$ for $t = 1 - 2\alpha$ or $t = 1 - \alpha$ is right admissible since $1 - 2\alpha = \gamma_b$ and $1 - \alpha = \gamma_c$. On the contrary, for $t = 2 - 3\alpha$, it is not right admissible because $t = T^{-1}(\gamma_c)$ but $\gamma_c < t$ contradicting (ii).

The following result is Theorem 14 in [24].

Theorem 17 (Rauzy) *Let T be a regular s -interval exchange transformation and let I be a right admissible interval for T . The transformation induced by T on I is a regular s -interval exchange transformation.*

Example 18 Consider again the transformation of Example 8. The transformation induced by T on the semi-interval $I = [0, 2\alpha[$ is the 3-interval exchange transformation represented in Figure 5.




Figure 5: The transformation induced on I .

The notion of left admissible interval is symmetrical to that of right admissible. For $\ell < t < r$, the semi-interval $[t, r[$ is *left admissible* for T if there is a $k \in \mathbb{Z}$ such that $t = T^k(\gamma_a)$ for some $a \in A$ and

- (i) if $k > 0$, then $T^h(\gamma_a) < t$ for all h such that $0 < h < k$,
- (ii) if $k \leq 0$, then $T^h(\gamma_a) < t$ for all h such that $k < h \leq 0$.

We also say that t itself is left admissible. Note that, as for right induction, the semi-intervals $[\gamma_a, r[$ and $[\nu_a, r[$ are left admissible. The symmetrical statements of Theorem 17 also hold for left admissible intervals.

Let now generalize the notion of admissibility to a two-sided version. For a semi-interval $I = [u, v[\subset [\ell, r[$, we define the following functions on $[\ell, r[$:

$$\rho_{I,T}^+(z) = \min\{n > 0 \mid T^n(z) \in]u, v]\}, \quad \rho_{I,T}^-(z) = \min\{n \geq 0 \mid T^{-n}(z) \in]u, v]\}.$$

We then define three sets. First, let

$$E_{I,T}(z) = \{k \mid -\rho_{I,T}^-(z) \leq k < \rho_{I,T}^+(z)\}.$$

Next, the set of *neighbors* of z with respect to I and T is

$$N_{I,T}(z) = \{T^k(z) \mid k \in E_{I,T}(z)\}.$$

The set of *division points* of I with respect to T is the finite set

$$\text{Div}(I, T) = \bigcup_{i=1}^s N_{I,T}(\gamma_i).$$

We now formulate the following definition. For $\ell \leq u < v \leq r$, we say that the semi-interval $I = [u, v[$ is *admissible* for T if $u, v \in \text{Div}(I, T) \cup \{r\}$.

Note that a semi-interval $[\ell, v[$ is right admissible if and only if it is admissible and that a semi-interval $[u, r[$ is left admissible if and only if it is admissible. Note also that $[\ell, r[$ is admissible.

Note also that for a regular interval exchange transformation relative to a partition $(I_a)_{a \in A}$, each of the semi-intervals I_a (or J_a) is admissible although only the first one is right admissible (and the last one is left admissible). Actually, we can prove that for every word w , the semi-intervals I_w and J_w are admissible. In order to do that, we need the following Lemma.

Lemma 19 *Let T be a s -interval exchange transformation on the semi-interval $[\ell, r[$. For any $k \geq 1$, the set $P_k = \{T^h(\gamma_i) \mid 1 \leq i \leq s, 1 \leq h \leq k\}$ is the set of $(s-1)k+1$ left boundaries of the semi-intervals J_y for all words $y \in F(T) \cap A^k$.*

Proof. Let Q_k be the set of left boundaries of the intervals J_y for $|y| = k$. Since $\text{Card}(F(T) \cap A^k) = (s-1)k+1$ by Proposition 10, we have $\text{Card}(Q_k) = (s-1)k+1$. Since T is regular the set $R_k = \{T^h(\gamma_i) \mid 2 \leq i \leq s, 1 \leq h \leq k\}$ is made of $(s-1)k$ distinct points. Moreover, since

$$\gamma_1 = T(\gamma_{\pi(1)}), T(\gamma_1) = T^2(\gamma_{\pi(1)}), \dots, T^{k-1}(\gamma_1) = T^k(\gamma_{\pi(1)}),$$

we have $P_k = R_k \cup \{T^k(\gamma_1)\}$. This implies $\text{Card}(P_k) \leq (s-1)k+1$. On the other hand, if $y = b_0 \cdots b_{k-1}$, then $J_y = \cap_{i=0}^{k-1} T^{k-i}(I_{b_i})$. Thus the left boundary of each J_y is the left boundary of some $T^h(I_a)$ for some h with $1 \leq h \leq k$ and some $a \in A$. Consequently $Q_k \subset P_k$. This proves that $\text{Card}(P_k) = (s-1)k+1$ and that consequently $P_k = Q_k$. ■

A dual statement holds for the semi-intervals I_y .

Proposition 20 *Let T be a s -interval exchange transformation on the semi-interval $[\ell, r[$. For any $w \in F(T)$, the semi-interval J_w is admissible.*

Proof. Set $|w| = k$ and $J_w = [u, v[$. By Lemma 19, we have $u = T^g(\gamma_i)$ for $1 \leq i \leq s$ and $1 \leq g \leq k$. Similarly, we have $v = r$ or $v = T^d(\gamma_j)$ for $1 \leq j \leq s$ and $1 \leq d \leq k$.

For $1 < h < g$, the point $T^h(\gamma_i)$ is the left boundary of some semi-interval J_y with $|y| = k$ and thus $T^h(\gamma_i) \notin J_w$. This shows that $g \in E_{J_w, T}(\gamma_i)$ and thus that $u \in \text{Div}(J_w, T)$.

If $v = r$, then $v \in \text{Div}(J_w, T)$. Otherwise, one shows in the same way as above that $v \in \text{Div}(J_w, T)$. Thus J_w is admissible. ■

Note that the same statement holds for the semi-intervals I_w instead of the semi-intervals J_w (using the dual statement of Lemma 19).

It can be useful to reformulate the definition of a division point and of an admissible pair using the terminology of graphs. Let $G(T)$ be the graph with vertex set $[\ell, r[$ and edges the pairs $(z, T(z))$ for $z \in [\ell, r[$. Then, if T is minimal and I is a semi-interval, for any $z \in [\ell, r[$, there is a path $P_{I, T}(z)$ such that its origin x and its end y are in I , z is on the path, $z \neq y$ and no vertex of the path except x, y are in I (actually $x = T^{-n}(z)$ with $n = \rho_{I, T}^-(z)$ and $y = T^m(z)$ with $m = \rho_{I, T}^+(z)$). Then the division points of I are the vertices which are on a path $P_{I, T}(\gamma_i)$ but not at its end (see Figure 6).




Figure 6: The neighbors of z with respect to $I = [u, v[$.

The following is a generalization of Theorem 17. Recall that $\text{Sep}(T)$ denotes the set of separation points of T , i.e. the points $\gamma_1 = 0, \gamma_2, \dots, \gamma_s$ (which are the left boundaries of the semi-intervals I_1, \dots, I_s).

Theorem 21 *Let T be a regular s -interval exchange transformation on $[\ell, r[$. For any admissible semi-interval $I = [u, v[$, the transformation S induced by T on I is a regular s -interval exchange transformation with separation points $\text{Sep}(S) = \text{Div}(I, T) \cap I$.*

Proof. Since T is regular, it is minimal. Thus for each $i \in \{2, \dots, s\}$ there are points $x_i, y_i \in]u, v[$ such that there is a path from x_i to y_i passing by γ_i but not containing any point of I except at its origin and its end. Since T is regular, the x_i are all distinct and the y_i are all distinct.

Since I is admissible, there exist $g, d \in \{1, \dots, s\}$ such that $u \in N_{I,T}(\gamma_g)$ and $v \in N_{I,T}(\gamma_d)$. Moreover, since u is a neighbor of γ_g with respect to I , u is on the path from x_g to y_g (it can be either before or after γ_g). Similarly, v is on the path from x_d to y_d (see Figure 7 where u is before γ_g and v is after γ_d).




Figure 7: The transformation induced on $[u, v[$.

Set $x_1 = y_1 = u$. Let $(I_j)_{1 \leq j \leq s}$ be the partition of I in semi-intervals such that x_j is the left boundary of I_j for $1 \leq j \leq s$. Let J_j be the partition of I such that y_j is the left boundary of J_j for $1 \leq j \leq s$. We will prove that

$$S(I_j) = \begin{cases} J_j & \text{if } j \neq 1, g \\ J_1 & \text{if } j = g \\ J_g & \text{if } j = 1 \end{cases}$$

and that the restriction of S to I_j is a translation.

Assume first that $j \neq 1, g$. Then $S(x_j) = y_j$. Let k be such that $y_j = T^k(x_j)$ and denote $I'_j = I_j \setminus x_j$. We will prove by induction on h that for $0 \leq h \leq k-1$, the set $T^h(I'_j)$ does not contain u, v or any x_i . It is true for $h = 0$. Assume that it holds up to $h < k-1$.

For any h' with $0 \leq h' \leq h$, the set $T^{h'}(I'_j)$ does not contain any γ_i . Indeed, otherwise there would exist h'' with $0 \leq h'' \leq h'$ such that $x_i \in T^{h''}(I'_j)$, a contradiction. Thus T is a translation on $T^{h'}(I'_j)$. This implies that T^h is a translation on I_j . Note also that $T^h(I'_j) \cap I = \emptyset$. Assume the contrary. We first observe that we cannot have $T^h(x_j) \in I$. Indeed, $h < k$ implies that $T^h(x_j) \notin]u, v[$. And we cannot have $T^h(x_j) = u$ since $j \neq g$. Thus $T^h(I'_j) \cap I \neq \emptyset$ implies that $u \in T^h(I'_j)$, a contradiction.

Suppose that $u = T^{h+1}(z)$ for some $z \in I'_j$. Since u is on the path from x_g to y_g , it implies that for some h' with $0 \leq h' \leq h$ we have $x_g = T^{h'}(z)$, a contradiction with the induction hypothesis. A similar proof (using the fact that v is on the path from x_d to y_d) shows that $T^{h+1}(I'_j)$ does not contain v . Finally suppose that some x_i is in $T^{h+1}(I'_j)$. Since the restriction of T^h to I_j is a translation, $T^h(I_j)$ is a semi-interval. Since $T^{h+1}(x_j)$ is not in I the fact that $T^{h+1}(I'_j) \cap I$ is not empty implies that $u \in T^h(I_j)$, a contradiction.

This shows that T^k is continuous at each point of I'_j and that $S = T^k(x)$ for all $x \in I_j$. This implies that the restriction of S to I_j is a translation into J_j .

If $j = 1$, then $S(x_1) = S(u) = y_g$. The same argument as above proves that the restriction of S to I_1 is a translation from I_1 into J_g . Finally if $j = g$, then $S(x_g) = x_1 = u$ and, similarly, we obtain that the restriction of S to I_g is a translation into I_1 .

Since S is the transformation induced by the transformation T which is one to one, it is also one to one. This implies that the restriction of S to each of the semi-intervals I_j is a bijection onto the corresponding interval J_j, J_1 or J_g according to the value of j .

This shows that S is an s -interval exchange transformation. Since the orbits of the points x_2, \dots, x_s relative to S are included in the orbits of $\gamma_2, \dots, \gamma_s$, they are infinite and disjoint. Thus S is regular.

Let us finally show that $\text{Sep}(S) = \text{Div}(I, T) \cap I$. We have $\text{Sep}(S) = \{x_1, x_2, \dots, x_s\}$ and $x_i \in N_{I, T}(\gamma_i)$. Thus $\text{Sep}(S) \subset \text{Div}(I, T) \cap I$. Conversely, let $x \in \text{Div}(I, T) \cap I$. Then $x \in N_{I, T}(\gamma_i) \cap I$ for some $1 \leq i \leq s$. If $i \neq 1, g$, then $x = x_i$. If $i = 1$, then either $x = u$ (if $u = \ell$) or $x = x_{\pi(1)}$ since $\gamma_1 = T(\gamma_{\pi(1)})$. Finally, if $i = g$ then $x = u$ or $x = x_g$. Thus $x \in \text{Sep}(S)$ in all cases. ■

Note that for any s -interval exchange transformation on $[\ell, r[$ and any semi-interval I of $[\ell, r[$, the transformation S induced by T on I is an interval exchange transformation on at most $s+2$ -intervals (see [12], Chapter 5 p. 128). Actually, it follows from the proof of Lemma 2, page 128 in [12] that, if T is regular and S is an s -interval exchange transformation with separation points $\text{Sep}(S) = \text{Div}(I, T) \cap I$, then I is admissible. Thus the converse of Theorem 21 is also true.

2.4 A closure property

In the following we will prove a closure property of the family of regular interval exchange sets. The same property holds for Sturmian sets (see [18]) and for uniformly recurrent tree sets (see [5]).

Lemma 22 *Let T be a regular interval exchange transformation and let $F = F(T)$. For $w \in F$, let S be the transformation induced by T on J_w . One has $x \in \mathcal{R}_F(w)$ if and only if*

$$\Sigma_T(z) = x\Sigma_T(S(z))$$

for some $z \in J_w$.

Proof. Assume first that $x \in \mathcal{R}_F(w)$. Then for any $z \in J_w \cap I_x$, we have $S(z) = T^{|x|}(z)$ and

$$\Sigma_T(z) = x\Sigma_T(T^{|x|}(z)) = x\Sigma_T(S(z)).$$

Conversely, assume that $\Sigma_T(z) = x\Sigma_T(S(z))$ for some $z \in J_w$. Then $T^{|x|}(z) \in J_w$ and thus $wx \in A^*w$ which implies that $x \in \Gamma_F(w)$. Moreover x does not have a proper prefix in $\Gamma_F(w)$ and thus $x \in \mathcal{R}_F(w)$. ■

Since a regular interval exchange set is recurrent, the previous lemma says that the natural coding of a point in J_w is a concatenation of first return words to w . Moreover, note also that $T^n(z) \in J_w$ if and only if the prefix of length n of $\Sigma_T(z)$ is a return word to w .

Theorem 23 *Any derived set of a regular s -interval exchange set is a regular s -interval exchange set.*

Proof. Let T be a regular s -interval exchange transformation and let $F = F(T)$.

Let $w \in F$. Since the semi-interval J_w is admissible according to Proposition 20, the transformation S induced by T on J_w is, by Theorem 21, an s -interval exchange transformation. The corresponding partition of J_w is the family $(J_{wx})_{w \in \mathcal{R}_F(w)}$.

Using Lemma 22 and the observation following, it is clear that $\Sigma_T(z) = f(\Sigma_S(z))$, where z is a point of J_w and $f : A^* \rightarrow \mathcal{R}_F(w)^*$ is a coding morphism for $\mathcal{R}_F(w)$.

Set $x = \Sigma_T(T^{-|w|}(z))$ and $y = \Sigma_T(z)$. Then $x = wy$ and thus $\Sigma_S(z) = \mathcal{D}_f(x)$. By Proposition 3, this shows that the derived set of F with respect to f is $F(S)$. ■

Theorem 23 implies, in particular, a result of [27], i.e., that $\text{Card}(\mathcal{R}_F(w)) = \text{Card}(A)$ (see also [3] and [5]).

3 Rauzy induction

In this section we describe the transformation called Rauzy induction defined in [24] which operates on regular interval transformations and recall the results concerning this transformation (Theorems 17 and 24). We introduce the definition of admissibility for an interval. It generalizes in a natural way the notion of admissibility defined in [24]. We also introduce a branching version of this transformation and generalize Rauzy's results to the two-sided case (Theorems 21 and 26). In particular we characterize in Theorem 26 the admissible semi-intervals for an interval exchange transformation.

3.1 One-sided Rauzy induction

Let $T = T_{\lambda, \pi}$ be a regular s -interval exchange transformation on $[\ell, r[$. Set $Z(T) = [\ell, \max\{\gamma_s, \delta_{\pi(s)}\}[$.

Note that $Z(T)$ is the largest semi-interval which is right-admissible for T . We denote by $\psi(T)$ the transformation induced by T on $Z(T)$.

The following result is Theorem 23 in [24].

Theorem 24 (Rauzy) *Let T be a regular interval exchange transformation. A semi-interval I is right admissible for T if and only if there is an integer $n \geq 0$ such that $I = Z(\psi^n(T))$. In this case, the transformation induced by T on I is $\psi^{n+1}(T)$.*

The map $T \mapsto \psi(T)$ is called the *right Rauzy induction*. There are actually two cases according to $\gamma_s < \delta_{\pi(s)}$ (Case 0) or $\gamma_s > \delta_{\pi(s)}$ (Case 1). We cannot have $\gamma_s = \delta_{\pi(s)}$ since T is regular.

In Case 0, we have $Z(T) = [\ell, \delta_{\pi(s)}[$ and for any $z \in Z(T)$,

$$\psi(T)(z) = \begin{cases} T^2(z) & \text{if } z \in I_{a_{\pi(s)}} \\ T(z) & \text{otherwise.} \end{cases}$$

The transformation $S = \psi(T)$ is the interval exchange transformation relative to $(K_a)_{a \in A}$ with $K_a = I_a \cap Z(T)$ for all $a \in A$. Note that $K_a = I_a$ for $a \neq a_s$. The translation values β_a are defined as follows, denoting α_i, β_i instead of $\alpha_{a_i}, \beta_{a_i}$,

$$\beta_i = \begin{cases} \alpha_{\pi(s)} + \alpha_s & \text{if } i = \pi(s) \\ \alpha_i & \text{otherwise.} \end{cases}$$

In summary, in Case 0, the semi-interval $J_{a_{\pi(s)}}$ is suppressed, the semi-interval J_{a_s} is split into $S(K_{a_s})$ and $S(K_{a_{\pi(s)}})$. The left boundaries of the semi-intervals K_a are the left boundaries of the semi-intervals I_a . The transformation is represented in Figure 8, in which the left boundary of the semi-interval $S(K_{a_{\pi(s)}})$ is denoted $\delta'_{\pi(s)}$.

In Case 1, we have $Z(T) = [\ell, \gamma_s[$ and for any $z \in Z(T)$,

$$\psi(T)(z) = \begin{cases} T^2(z) & \text{if } z \in T^{-1}(I_{a_s}) \\ T(z) & \text{otherwise.} \end{cases}$$




Figure 8: Case 0 in Rauzy induction.

The transformation $S = \psi(T)$ is the interval exchange transformation relative to $(K_a)_{a \in A}$ with

$$K_a = \begin{cases} T^{-1}(I_a) & \text{if } a = a_s \\ T^{-1}(T(I_a) \cap Z(T)) & \text{otherwise.} \end{cases}$$

Note that $K_a = I_a$ for $a \neq a_s$ and $a \neq a_{\pi(s)}$. Moreover $K_a = S^{-1}(T(I_a) \cap Z(T))$ in all cases. The translation values β_i are defined by

$$\beta_i = \begin{cases} \alpha_{\pi(s)} + \alpha_s & \text{if } i = s \\ \alpha_i & \text{otherwise.} \end{cases}$$

In summary, in Case 1, the semi-interval I_{a_s} is suppressed, the semi-interval $I_{a_{\pi(s)}}$ is split into $K_{a_{\pi(s)}}$ and K_{a_s} . The left boundaries of the semi-intervals $S(K_a)$ are the left boundaries of the semi-intervals J_a . The transformation is represented in Figure 9, where the left boundary of the semi-interval K_{a_s} is denoted γ'_s .




Figure 9: Case 1 in Rauzy induction.

Example 25 Consider again the transformation T of Example 8. Since $Z(T) = [0, 2\alpha[$, the transformation $\psi(T)$ is represented in Figure 5. The transformation $\psi^2(T)$ is represented in Figure 10.




Figure 10: The transformation $\psi^2(T)$.

The symmetrical notion of *left Rauzy induction* is defined similarly.

Let $T = T_{\lambda, \pi}$ be a regular s -interval exchange transformation on $[\ell, r[$. Set $Y(T) = [\min\{\mu_1, \nu_{\pi(1)}\}, r[$. We denote by $\varphi(T)$ the transformation induced by T on $Y(T)$. The map $T \mapsto \varphi(T)$ is called the *left Rauzy induction*.

Note that one has also $Y(T) = [\min\{\gamma_2, \delta_{\pi(2)}\}, r[$.

The symmetrical statements of Theorem 24 also hold for left admissible intervals.

3.2 Branching induction

The following is a generalization of Theorem 24.

Theorem 26 *Let T be a regular s -interval exchange transformation on $[\ell, r[$. A semi-interval I is admissible for T if and only if there is a sequence $\chi \in \{\varphi, \psi\}^*$ such that I is the domain of $\chi(T)$. In this case, the transformation induced by T on I is $\chi(T)$.*

We first prove the following lemmas, in which we assume that T is a regular s -interval exchange transformation on $[\ell, r[$. Recall that $Y(T), Z(T)$ are the domains of $\varphi(T), \psi(T)$ respectively.

Lemma 27 *If a semi-interval I strictly included in $[\ell, r[$ is admissible for T , then either $I \subset Y(T)$ or $I \subset Z(T)$.*

Proof. Set $I = [u, v[$. Since I is strictly included in $[\ell, r[$, we have either $\ell < u$ or $v < r$. Set $Y(T) = [y, r[$ and $Z(T) = [\ell, z[$.

Assume that $v < r$. If $y \leq u$, then $I \subset Y(T)$. Otherwise, let us show that $v \leq z$. Assume the contrary. Since I is admissible, we have $v = T^k(\gamma_i)$ with $k \in E_{I, T}(\gamma_i)$ for some i with $1 \leq i \leq s$. But $k > 0$ is impossible since $u < T(\gamma_i) < v$ implies $T(\gamma_i) \in]u, v[$, in contradiction with the fact that $k < \rho_I^+(\gamma_i)$. Similarly, $k \leq 0$ is impossible since $u < \gamma_i < v$ implies $\gamma_i \in]u, v[$. Thus $I \subset Z(T)$.

The proof in the case $\ell < u$ is symmetric. ■

The next lemma is the two-sided version of Lemma 22 in [24].

Lemma 28 *Let T be a regular s -interval exchange transformation on $[\ell, r[$. Let J be an admissible semi-interval for T and let S be the transformation induced by T on J . A semi-interval $I \subset J$ is admissible for T if and only if it is admissible for S . Moreover $\text{Div}(J, T) \subset \text{Div}(I, T)$.*

Proof. Set $J = [t, w[$ and $I = [u, v[$. Since J is admissible for T , the transformation S is a regular s -interval exchange transformation by Theorem 21.

Suppose first that I is admissible for T . Then $u = T^g(\gamma_i)$ with $g \in E_{I,T}(\gamma_i)$ for some $1 \leq i \leq s$, and $v = T^d(\gamma_j)$ with $d \in E_{I,T}(\gamma_j)$ for some $1 \leq j \leq s$ or $v = r$.

Since S is the transformation induced by T on J there is a separation point x of S of the form $x = T^m(\gamma_i)$ with $m = -\rho_{J,T}^-(\gamma_i)$ and thus $m \in E_{J,T}(\gamma_i)$. Thus $u = T^{g-m}(x)$.

Assume first that $g - m > 0$. Since $u, x \in J$, there is an integer n with $0 < n \leq g - m$ such that $u = S^n(x)$.

Let us show that $n \in E_{I,S}(x)$. Assume by contradiction that $\rho_{I,S}^+(x) \leq n$. Then there is some k with $0 < k \leq n$ such that $S^k(x) \in]u, v[$. But we cannot have $k = n$ since $u \notin]u, v[$. Thus $k < n$.

Next, there is h with $0 < h < g - m$ such that $T^h(x) = S^k(x)$. Indeed, setting $y = S^k(x)$, we have $u = T^{g-m-h}(y) = S^{n-k}(y)$ and thus $h < g - m$. If $0 < h \leq -m$, then $T^h(x) = T^{m+h}(\gamma_i) \in I \subset J$ contradicting the hypothesis that $m \in E_{J,T}(\gamma_i)$. If $-m < h < g - m$, then $T^h(x) = T^{m+h}(\gamma_i) \in I$, contradicting the fact that $g \in E_{I,T}(\gamma_i)$. This shows that $n \in E_{I,S}(x)$ and thus that $u \in \text{Div}(I, S)$.

Assume next that $g - m \leq 0$. There is an integer n with $g - m \leq n \leq 0$ such that $u = S^n(x)$. Let us show that $n \in E_{I,S}(x)$. Assume by contradiction that $n < -\rho_{I,S}^-(x)$. Then there is some k with $n < k < 0$ such that $S^k(x) = T^h(x)$. Then $T^h(x) = T^{h+m}(\gamma_i) \in I$ with $g < h + m < m$, in contradiction with the hypothesis that $m \in E_{I,T}(\gamma_i)$.

We have proved that $u \in \text{Div}(I, S)$. If $v = r$, the proof that I is admissible for S is complete. Otherwise, the proof that $v \in \text{Div}(I, S)$ is similar to the proof for u .

Conversely, if I is admissible for S , there is some $x \in \text{Sep}(S)$ and $g \in E_{I,S}(x)$ such that $u = S^g(x)$. But $x = T^m(\gamma_i)$ and since $u, x \in J$ there is some n such that $u = T^n(\gamma_i)$.

Assume for instance that $n > 0$ and suppose that there exists k with $0 < k < n$ such that $T^k(\gamma_i) \in]u, v[$. Then, since $I \subset J$, $T^k(\gamma_i)$ is of the form $S^h(x)$ with $0 < h < g$ which contradicts the fact that $g \in E_{I,S}(x)$. Thus $n \in E_{I,T}(\gamma_i)$ and $u \in \text{Div}(I, T)$.

The proof is similar in the case $n \leq 0$.

If $v = r$, we have proved that I is admissible for T . Otherwise, the proof that $v \in \text{Div}(I, T)$ is similar.

Finally, assume that I is admissible for T (and thus for S). For any $\gamma_i \in \text{Sep}(T)$, one has

$$\rho_{I,T}^-(\gamma_i) \geq \rho_{J,T}^-(\gamma_i) \quad \text{and} \quad \rho_{I,T}^+(\gamma_i) \geq \rho_{J,T}^+(\gamma_i)$$

showing that $\text{Div}(J, T) \subset \text{Div}(I, T)$. ■

The last lemma is the key argument to prove Theorem 26. It is a tree version of the argument used by Rauzy in [24].

Lemma 29 *For any admissible interval $I \subset [\ell, r[$, the set \mathcal{F} of sequences $\chi \in \{\varphi, \psi\}^*$ such that $I \subset D(\chi(T))$ is finite.*

Proof. The set \mathcal{F} is suffix-closed. Indeed it contains the empty word because $[\ell, r[$ is admissible. Moreover, for any $\xi, \chi \in \{\varphi, \psi\}^*$, one has $D(\xi\chi(T)) \subset D(\chi(T))$ and thus $\xi\chi \in \mathcal{F}$ implies $\chi \in \mathcal{F}$.

The set \mathcal{F} is finite. Indeed, by Lemma 28, applied to $J = D(\chi(T))$, for any $\chi \in \mathcal{F}$, one has $\text{Div}(D(\chi(T)), T) \subset \text{Div}(I, T)$. In particular, the boundaries of $D(\chi(T))$ belong to $\text{Div}(I, T)$. Since $\text{Div}(I, T)$ is a finite set, this implies that there is a finite number of possible semi-intervals $D(\chi(T))$. Thus there is no infinite word with all its suffixes in \mathcal{F} . Since the sequences χ are binary, this implies that \mathcal{F} is finite. ■

Proof of Theorem 26. We first prove by induction on the length of χ that the domain I of $\chi(T)$ is admissible and that the transformation induced by T on I is $\chi(T)$. It is true for $|\chi| = 0$ since $[\ell, r[$ is admissible and $\chi(T) = T$. Next, assume that $J = D(\chi(T))$ is admissible and that the transformation induced by T on J is $\chi(T)$. Then $D(\varphi\chi(T))$ is admissible for $\chi(T)$ since $D(\varphi\chi(T)) = Y(\chi(T))$. Thus $I = D(\varphi\chi(T))$ is admissible for T by Lemma 28 and the transformation induced by T on I is $\varphi\chi(T)$. The same proof holds for $\psi\chi$.

Conversely, assume that I is admissible. By Lemma 29, the set \mathcal{F} of sequences $\chi \in \{\varphi, \psi\}^*$ such that $I \subset D(\chi(T))$ is finite.

Thus there is some $\chi \in \mathcal{F}$ such that $\varphi\chi, \psi\chi \notin \mathcal{F}$. If I is strictly included in $D(\chi(T))$, then by Lemma 27 applied to $\chi(T)$, we have $I \subset Y(\chi(T)) = D(\varphi\chi(T))$ or $I \subset Z(\chi(T)) = D(\psi\chi(T))$, a contradiction. Thus $I = D(\chi(T))$. ■

We close this subsection with a result concerning the dynamics of the branching induction.

Theorem 30 *For any sequence $(T_n)_{n \geq 0}$ of regular interval exchange transformations such that $T_{n+1} = \varphi(T_n)$ or $T_{n+1} = \psi(T_n)$ for all $n \geq 0$, the length of the domain of T_n tends to 0 when $n \rightarrow \infty$.*

Proof. Assume the contrary and let I be an open interval included in the domain of T_n for all $n \geq 0$. The set $\text{Div}(I, T) \cap I$ is formed of s points. For any pair u, v of consecutive elements of this set, the semi-interval $[u, v[$ is admissible. By Lemma 29, there is an integer n such that the domain of T_n does not contain $[u, v[$, a contradiction. ■

3.3 Equivalence relation

Let $[\ell_1, r_1[$, $[\ell_2, r_2[$ be two semi-intervals of the real line. Let $T_1 = T_{\lambda, \pi_1}$ be an s -interval exchange transformation relative to a partition of $[\ell_1, r_1[$ and $T_2 = T_{\mu, \pi_2}$ another s -interval exchange transformations relative to $[\ell_2, r_2[$. We say that T_1 and T_2 are *equivalent* if $\pi_1 = \pi_2$ and $\lambda = c\mu$ for some $c > 0$. Thus, two interval exchange transformations are equivalent if we can obtain the second

from the first by a rescaling following by a translation. We denote by $[T_{\lambda,\pi}]$ the equivalence class of $T_{\lambda,\pi}$.

Example 31 Let $S = T_{\mu,\pi}$ be the 3-interval exchange transformation on a partition of the semi-interval $[2\alpha, 1[$, with $\alpha = (3 - \sqrt{5})/2$, represented in Figure 11. S is equivalent to the transformation $T = T_{\lambda,\pi}$ of Example 8, with length vector $\lambda = (1 - 2\alpha, \alpha, \alpha)$ and permutation the cycle $\pi = (132)$. Indeed the length vector $\mu = (8\alpha - 3, 2 - 5\alpha, 2 - 5\alpha)$ satisfies $\mu = \frac{2-5\alpha}{\alpha}\lambda$.




Figure 11: The transformation S .

Note that if T is a minimal (resp. regular) interval exchange transformation and $[S] = [T]$, then S is also minimal (resp. regular).

For an interval exchange transformation T we consider the directed labeled graph $\mathcal{G}(T)$, called the *induction graph* of T , defined as follows. The vertices are the equivalence classes of transformations obtained starting from T and applying all possible $\chi \in \{\psi, \varphi\}^*$. There is an edge labeled ψ (resp. φ) from a vertex $[S]$ to a vertex $[U]$ if and only if $U = \psi(S)$ (resp. $\varphi(S)$) for two transformations $S \in [S]$ and $U \in [U]$.

Example 32 Let $\alpha = \frac{3-\sqrt{5}}{2}$ and R be a rotation of angle α . By Example 5, R is a 2-interval exchange transformation on $[0, 1[$ relative to the partition $[0, 1 - \alpha[$, $[1 - \alpha, 1[$. The induction graph $\mathcal{G}(R)$ of the transformation is represented in the left of Figure 13.

Note that for a 2-interval exchange transformation T , one has $[\psi(T)] = [\varphi(T)]$, whereas in general the two transformations are not equivalent.

The induction graph of an interval exchange transformation can be infinite. A sufficient condition for the induction graph to be finite is given in Section 4.

Let now introduce a variant of this equivalence relation (and of the related graph). We consider the case of two transformations “equivalent” up to reflection (and up to the separation points). This choice allows us to obtain the same natural coding for an interval exchange transformation relative to a point, and for the mirror transformation relative to the specular point (with respect to the midpoint of the of the interval).

For an s -interval exchange transformation $T = T_{\lambda,\pi}$, with length vector $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_s)$, we define the *mirror transformation* $\tilde{T} = T_{\tilde{\lambda},\tau\circ\pi}$ of T , where $\tilde{\lambda} = (\lambda_s, \lambda_{s-1}, \dots, \lambda_1)$ and $\tau : i \mapsto (s - 1 + 1)$ is the permutation that reverses the names of the semi-intervals.

Given two interval exchange transformations T_1 and T_2 on the same alphabet relative to two partitions of two semi-intervals $[\ell_1, r_1[$ and $[\ell_2, r_2[$ respectively, we say that T_1 and T_2 are *similar* either if $[T_1] = [T_2]$ or $[T_1] = [\widetilde{T}_2]$. Clearly, similarity is also an equivalent relation. We denote by $\langle T \rangle$ the class of transformations similar to T .

Example 33 Let T be the interval exchange transformation of Example 8. The transformation $U = \varphi^6(T)$ is represented in Figure 12 (see also Example 41). It is easy to verify that U is similar to the transformation S of Example 31. Indeed, we can obtain the second transformation (up to the separation points and the end points) by taking the mirror image of the domain.

Note that the order of the labels, i.e. the order of the letters of the alphabet, may be different from the order of the original transformation.




Figure 12: The transformation U .

As of the equivalence relation, also similarity preserves minimality and regularity.

Let T be an interval exchange transformation. We denote by

$$\mathcal{S}(T) = \bigcup_{n \in \mathbb{Z}} T^n(\text{Sep}(T))$$

the union of the orbits of the separation points. Let S be an interval exchange transformation similar to T . Thus, there exists a bijection $f : D(T) \setminus \mathcal{S}(T) \rightarrow D(S) \setminus \mathcal{S}(S)$. This bijection is given by an affine transformation, namely a rescaling following by a translation if T and S are equivalent and a rescaling following by a translation and a reflection otherwise. By the previous remark, if T is a minimal exchange interval transformation and S is similar to T , then the two interval exchange sets $F(T)$ and $F(S)$ are equal up to permutation, that is there exists a permutation π such that one for every $w = a_0 a_1 \cdots a_{n-1} \in F(T)$ there exists a unique word $v = b_0 b_1 \cdots b_{n-1} \in F(S)$ such that $b_i = \pi(a_i)$ for all $i = 1, 2, \dots, n-1$.

In a similar way as before, we can use the similarity in order to construct a graph. For an interval exchange transformation T we define $\tilde{\mathcal{G}}(T)$ the *modified induction graph* of T as the directed (unlabeled) graph with vertices the similar classes of transformations obtained starting from T and applying all possible $\chi \in \{\psi, \varphi\}^*$ and an edge from $\langle S \rangle$ to $\langle U \rangle$ if $U = \psi(S)$ or $U = \varphi(S)$ for two transformations $S \in \langle S \rangle$ and $U \in \langle U \rangle$.

Note that this variant appears naturally when considering the Rauzy induction of a 2-interval exchange transformation as a continued fraction expansion. There exists a natural bijection between the closed interval $[0, 1]$ of the real line and the set of 2-interval exchange transformation given by the map $x \mapsto T_{\lambda, \pi}$ where $\pi = (12)$ and $\lambda = (\lambda_1, \lambda_2)$ is the length vector such that $x = \frac{\lambda_1}{\lambda_2}$.

In this view, the Rauzy induction corresponds to the Euclidean algorithm (see [22] for more details), i.e. the map $\mathcal{E} : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+^2$ given by

$$\mathcal{E}(\lambda_1, \lambda_2) = \begin{cases} (\lambda_1 - \lambda_2, \lambda_2) & \text{if } \lambda_1 \geq \lambda_2 \\ (\lambda_1, \lambda_2 - \lambda_1) & \text{otherwise.} \end{cases}$$

Applying iteratively the Rauzy induction starting from T corresponds then to the continued fraction expansion of x .

Example 34 Let α and R be as in Example 32. The modified induction graph $\tilde{\mathcal{G}}(R)$ of the transformation is represented on the right of Figure 13. Note that the ratio of the two lengths of the semi-intervals exchanged by T is $\frac{1-\alpha}{\alpha} = \frac{1+\sqrt{5}}{2} = \phi = 1 + \frac{1}{1+\dots}$.



Figure 13: Induction graph and modified induction graph of the rotation R of angle $\alpha = (3 - \sqrt{5})/2$.

3.4 Induction and automorphisms

Let $T = T_{\lambda, \pi}$ be a regular interval exchange on $[\ell, r[$ relative to $(I_a)_{a \in A}$. Set $A = \{a_1, \dots, a_s\}$. Recall now from Subsection 2.2 that for any $z \in [\ell, r[$, the natural coding of T relative to z is the infinite word $\Sigma_T(z) = b_0 b_1 \dots$ on the alphabet A with $b_n \in A$ defined for $n \geq 0$ by $b_n = a$ if $T^n(z) \in I_a$.

Denote by θ_1, θ_2 the morphisms from A^* into itself defined by

$$\theta_1(a) = \begin{cases} a_{\pi(s)} a_s & \text{if } a = a_{\pi(s)} \\ a & \text{otherwise} \end{cases}, \quad \theta_2(a) = \begin{cases} a_{\pi(s)} a_s & \text{if } a = a_s \\ a & \text{otherwise} \end{cases}.$$

The morphisms θ_1, θ_2 extend to automorphisms of the free group on A .

The following result already appears in [17]. We give a proof for the sake of completeness.

Proposition 35 *Let T be a regular interval exchange transformation on the alphabet A and let $S = \psi(T)$, $I = Z(T)$. There exists an automorphism θ of the free group on A such that $\Sigma_T(z) = \theta(\Sigma_S(z))$ for any $z \in I$.*

Proof. Assume first that $\gamma_s < \delta_{\pi(s)}$ (Case 0). We have $Z(T) = [\ell, \delta_{\pi(s}[$ and for any $x \in Z(T)$,

$$S(z) = \begin{cases} T^2(z) & \text{if } z \in K_{a_{\pi(s)}} = I_{a_{\pi(s)}} \\ T(z) & \text{otherwise.} \end{cases}$$

We will prove by induction on the length of w that for any $z \in I$, $\Sigma_S(z) \in wA^*$ if and only if $\Sigma_T(z) \in \theta_1(w)A^*$. The property is true if w is the empty word. Assume next that $w = av$ with $a \in A$ and thus that $z \in I_a$. If $a \neq a_{\pi(s)}$, then $\theta_1(a) = a$, $S(z) = T(z)$ and

$$\Sigma_S(z) \in avA^* \Leftrightarrow \Sigma_S(S(z)) \in vA^* \Leftrightarrow \Sigma_T(T(z)) \in \theta_1(v)A^* \Leftrightarrow \Sigma_T(z) \in \theta_1(w)A^*.$$

Otherwise, $\theta_1(a) = a_{\pi(s)}a_s$, $S(z) = T^2(z)$. Moreover, $\Sigma_T(z) = a_{\pi(s)}a_s\Sigma_T(T^2(z))$ and thus

$$\Sigma_S(z) \in avA^* \Leftrightarrow \Sigma_S(S(z)) \in vA^* \Leftrightarrow \Sigma_T(T^2(z)) \in \theta_1(v)A^* \Leftrightarrow \Sigma_T(z) \in \theta_1(w)A^*.$$

If $\delta_{\pi(s)} < \gamma_s$ (Case 1), we have $Z(T) = [\ell, \gamma_s[$ and for any $z \in Z(T)$,

$$S(z) = \begin{cases} T^2(z) & \text{if } z \in K_{a_s} = T^{-1}(I_{a_s}) \\ T(z) & \text{otherwise.} \end{cases}$$

As in Case 0, we will prove by induction on the length of w that for any $z \in I$, $\Sigma_S(z) \in wA^*$ if and only if $\Sigma_T(z) \in \theta_2(w)A^*$.

The property is true if w is empty. Assume next that $w = av$ with $a \in A$. If $a \neq a_s$, then $\theta_2(a) = a$, $S(z) = T(z)$ and $z \in K_a \subset I_a$. Thus

$$\Sigma_S(z) \in avA^* \Leftrightarrow \Sigma_S(S(z)) \in vA^* \Leftrightarrow \Sigma_T(T(z)) \in \theta_2(v)A^* \Leftrightarrow \Sigma_T(z) \in \theta_2(w)A^*.$$

Next, if $a = a_s$, then $\theta_2(a) = a_{\pi(s)}a_s$, $S(z) = T^2(z)$ and $z \in K_{a_s} = T^{-1}(I_{a_s}) \subset I_{a_{\pi(s)}}$. Thus

$$\Sigma_S(z) \in avA^* \Leftrightarrow \Sigma_S(S(z)) \in vA^* \Leftrightarrow \Sigma_T(T^2(z)) \in \theta_2(v)A^* \Leftrightarrow \Sigma_T(z) \in \theta_2(w)A^*.$$

where the last equivalence results from the fact that $\Sigma_T(z) \in a_{\pi(s)}a_sA^*$. This proves that $\Sigma_T(z) = \theta_2(\Sigma_S(z))$. \blacksquare

Example 36 Let T be the transformation of Example 8. The automorphism θ_1 is defined by

$$\theta_1(a) = ac, \quad \theta_1(b) = b, \quad \theta_1(c) = c.$$

The right Rauzy induction gives the transformation $S = \psi(T)$ computed in Example 18. One has $\Sigma_S(\alpha) = bacba \cdots$ and $\Sigma_T(\alpha) = baccbac \cdots = \theta_1(\Sigma_S(\alpha))$.

We state the symmetrical version of Proposition 35 for left Rauzy induction. The proof is analogous.

Proposition 37 *Let T be a regular interval exchange transformation on the alphabet A and let $S = \varphi(T)$, $I = Y(T)$. There exists an automorphism θ of the free group on A such that $\Sigma_T(z) = \theta(\Sigma_S(z))$ for any $z \in I$.*

Combining Propositions 35 and 37, we obtain the following statement.

Theorem 38 *Let T be a regular interval exchange transformation. For $\chi \in \{\varphi, \psi\}^*$, let $S = \chi(T)$ and let I be the domain of S . There exists an automorphism θ of the free group on A such that $\Sigma_T(z) = \theta(\Sigma_S(z))$ for all $z \in I$.*

Proof. The proof follows easily by induction on the length of χ using Propositions 35 and 37. ■

Note that if the transformations T and $S = \chi(T)$, with $\chi \in \{\psi, \varphi\}^*$, are equivalent, then there exists a point $z_0 \in D(S) \subseteq D(T)$ such that z_0 is a fixed point of the isometry that transforms $D(S)$ into $D(T)$ (if χ is different from the identity map, this point is unique). In that case one has $\Sigma_S(z_0) = \Sigma_T(z_0) = \theta(\Sigma_S(z_0))$ for an appropriate automorphism θ , i.e. $\Sigma_T(z_0)$ is a fixed point of an appropriate automorphism.

Corollary 39 *Let T be a regular interval exchange transformation. For $w \in F(T)$, the set $\mathcal{R}_F(w)$ is a basis of the free group on A .*

Proof. By Proposition 20, the semi-interval J_w is admissible. By Theorem 26 there is a sequence $\chi \in \{\varphi, \psi\}^*$ such that $D(\chi(T)) = J_w$. Moreover, the transformation $S = \chi(T)$ is the transformation induced by T on J_w . By Theorem 38 there is an automorphism θ of the free group on A such that $\Sigma_T(z) = \theta(\Sigma_S(z))$ for any $z \in J_w$.

By Lemma 22, we have $x \in \mathcal{R}_F(w)$ if and only if $\Sigma_T(z) = x\Sigma_T(S(z))$ for some $z \in J_w$. This implies that $\mathcal{R}_F(w) = \theta(A)$. Indeed, for any $z \in J_w$, let a is the first letter of $\Sigma_S(z)$. Then

$$\Sigma_T(z) = \theta(\Sigma_S(z)) = \theta(a\Sigma_S(S(z))) = \theta(a)\theta(\Sigma_S(Sz)) = \theta(a)\Sigma_T(S(z)).$$

Thus $x \in \mathcal{R}_F(w)$ if and only if there is $a \in A$ such that $x = \theta(a)$. This proves that the set $\mathcal{R}_F(w)$ is a basis of the free group on A . ■

The property proved in the previous corollary is actually true for a much larger class of sets than regular interval exchange sets (see [5, Theorem 4.7]). We illustrate this result with the following examples.

Example 40 We consider again the transformation T of Example 8 and $F = F(T)$. We have $\mathcal{R}_F(c) = \{bac, bbac, c\}$ (see Example 14). We represent in Figure 14 the sequence χ of Rauzy inductions such that J_c is the domain of $\chi(T)$.

The sequence is composed of a right induction followed by two left inductions. We have indicated on each edge the associated automorphism (indicating only




Figure 14: The sequence $\chi \in \{\varphi, \psi\}^*$

the image of the letter which is modified). We have $\chi = \varphi^2\psi$ and the resulting composition θ of automorphisms gives

$$\theta(a) = bac, \quad \theta(b) = bbac, \quad \theta(c) = c.$$

Thus $\mathcal{R}_F(c) = \theta(A)$.

Example 41 Let T and F be as in the preceding example. Let U be the transformation induced by T on J_a . We have $U = \varphi^6(T)$ and a computation shows that for any $z \in J_a$, $\Sigma_T(z) = \theta(\Sigma_U(z))$ where θ is the automorphism of the free group on $A = \{a, b, c\}$ which is the coding morphism for $\mathcal{R}_F(a)$ defined by:

$$\theta(a) = ccba, \quad \theta(b) = cbba, \quad \theta(c) = ccbba.$$

One can verify that $F(U) = F(S)$, where S is the transformation obtain from T by permuting the labels of the intervals according to the permutation $\pi = (acb)$.

Note that $F(U) = F(S)$ although S and U are not identical, even up to rescaling the intervals. Actually, the rescaling of U to a transformation on $[0, 1[$ corresponds to the mirror image of S , obtained by taking the image of the intervals by a symmetry centered at $1/2$.

Note that in the above examples, all lengths of the intervals belong to the quadratic number field $\mathbb{Q}[\sqrt{5}]$.

In the next Section we will prove that if a regular interval exchange transformation T is defined over a quadratic field, then the family of transformations obtained from T by the Rauzy inductions contains finitely many distinct transformations up to rescaling.

4 Interval exchange over a quadratic field

An interval exchange transformation is said to be defined over a set $Q \subset \mathbb{R}$ if the lengths of all exchanged semi-intervals belong to Q .

The following is proved in [10]. Let T be a minimal interval exchange transformation on semi-intervals defined over a quadratic number field. Let $(T_n)_{n \geq 0}$

be a sequence of interval exchange transformation such that $T_0 = T$ and T_{n+1} is the transformation induced by T_n on one of its exchanged semi-intervals I_n . Then, up to rescaling all semi-intervals I_n to the same length, the sequence (T_n) contains finitely many distinct transformations. In the same paper, an extension to the right Rauzy induction is suggested (but not completely developed).

In this section we generalize this results and prove that, under the above hypothesis on the lengths of the semi-intervals and up to rescaling and translation, there are finitely many transformations obtained by the branching Rauzy induction defined in Section 3.

Theorem 42 *Let T be a regular interval exchange transformation defined over a quadratic field. The family of all induced transformation of T over an admissible semi-interval contains finitely many distinct transformations up to equivalence.*

The proof of the Theorem 42 is based on the fact that for each minimal interval exchange transformation defined over a quadratic field, a certain measure of the arithmetic complexity of the admissible semi-intervals is bounded.

4.1 Complexities

Let T be an interval exchange transformation on a semi-interval $[\ell, r[$ defined over a quadratic field $\mathbb{Q}[\sqrt{d}]$, where d is a square free integer ≥ 2 . Without loss of generality, one may assume, by replacing T by an equivalent interval exchange transformation if necessary, that T is defined over the ring $\mathbb{Z}[\sqrt{d}] = \{m + n\sqrt{d} \mid m, n \in \mathbb{Z}\}$ and that all γ_i and α_i lie in $\mathbb{Z}[\sqrt{d}]$ (replacing $[\ell, r[$ if necessary by its equivalent translate with $\gamma_0 = \ell \in \mathbb{Z}[\sqrt{d}]$).

For $z = m + n\sqrt{d}$, let define $\Psi(z) = \max(|m|, |n|)$.

Let $\mathcal{A}([\ell, r[)$ be the algebra of subsets $X \subset [\ell, r[$ which are finite unions $X = \bigcup_j I_j$ of semi-intervals defined over $\mathbb{Z}[\sqrt{d}]$, that is $I_j = [\ell_j, r_j[$ for some $\ell_j, r_j \in \mathbb{Z}[\sqrt{d}]$. Note that the algebra $\mathcal{A}([\ell, r[)$ is closed under taking finite unions, intersections and passing to complements in $[\ell, r[$.

Set $\partial(X)$ the boundary of X and $|X|$ the Lebesgue measure of X . Given a subset $X \in \mathcal{A}([\ell, r[)$, we define the *complexity* of X as $\Psi(X) = \max\{\Psi(z) \mid z \in \partial(X)\}$ and the *reduced complexity* of X as $\Pi(X) = |X| \Psi(X)$.

A key tool to prove Theorem 42 is the following result proved in [10, Theorem 3.1].

Theorem 43 (Boshernitzan) *Let T be a minimal interval exchange transformation on an interval $[\ell, r[$ defined over a quadratic number field. Assume that $(J_n)_{\geq 1}$ is a sequence of semi-intervals of $[\ell, r[$ such that the set $\{\Pi(J_n) \mid n \geq 1\}$ is bounded. Then the sequence T_n of interval exchange transformations obtained by inducing T on J_n contains finitely many distinct equivalence classes of interval exchange transformations.*

Thus, in order to prove Theorem 42, it is sufficient to show that the reduced complexity of every admissible semi-interval is bounded.

The following Proposition is proved in [10, Proposition 2.1]. It shows that the complexity of a subset X and of its image $T(X)$ differ at most by a constant that depends only on T .

Proposition 44 *There exists a constant $u = u(T)$ such that for every $X \in \mathcal{A}([\ell, r[)$ and $z \in [\ell, r[$ one has $|\Psi(T(X)) - \Psi(X)| \leq u$ and $\Psi(T(z) - z) \leq u$. Moreover, one has $\Psi(\gamma) \leq u$ and $\Psi(T(\gamma)) \leq u$ for every separation point γ .*

Clearly, by Proposition 44, one also has $|\Psi(T^{-1}(X)) - \Psi(X)| \leq u$ for every $X \in \mathcal{A}([\ell, r[)$ and $\Psi(T^{-1}(z) - z) \leq u$ for every $z \in [\ell, r[$.

Although it is not necessary for our purposes, we can improve the approximation of the reduced complexity of a nonempty subset $X \in \mathcal{A}([\ell, r[)$ by the following proposition. This result, proved in [10, Proposition 2.4], determines a lower bound on $\Pi(X)$.

Proposition 45 *Let $X \in \mathcal{A}([\ell, r[)$ be a subset composed of n disjoint semi-intervals. Then $\Pi(X) > n/(4\sqrt{d})$.*

4.2 Return times

Let T be an interval exchange transformation. For a subset $X \in \mathcal{A}([\ell, r[)$ we define the *maximal positive return time* and *maximal negative return time* for T on X by the functions

$$\rho^+(X) = \min \left\{ n \geq 1 \mid T^n(X) \subset \bigcup_{i=0}^{n-1} T^i(X) \right\},$$

and

$$\rho^-(X) = \min \left\{ m \geq 1 \mid T^m(X) \subset \bigcup_{i=0}^{m-1} T^{-i}(X) \right\}.$$

We also define the *minimal positive return time* and the *minimal negative return time* as

$$\sigma^+(X) = \min \{ n \geq 1 \mid T^n(X) \cap X \neq \emptyset \},$$

and

$$\sigma^-(X) = \min \{ m \geq 1 \mid T^{-m}(X) \cap X \neq \emptyset \}.$$

If T is minimal, it is clear that for every $X \in \mathcal{A}([\ell, r[)$, one has

$$[\ell, r[= \bigcup_{i=0}^{\rho^+(X)-1} T^i(X) = \bigcup_{i=0}^{\rho^-(X)-1} T^{-i}(X).$$

Note that when J is a semi-interval, we have $\rho^+(J) = \max_{z \in J} \rho_{J,T}^+(z)$ and $\sigma^+(J) = \min_{z \in J} \rho_{J,T}^+(z)$. Symmetrically $\rho^-(J) = \max_{z \in J} \rho_{J,T}^-(z) + 1$ and $\sigma^-(J) = \min_{z \in J} \rho_{J,T}^-(z) + 1$.

Let ζ, η be two functions. We write $\zeta \in O(\eta)$ if there exists a constant C such that $|\zeta| \leq C|\eta|$. We write $\zeta \in \Theta(\eta)$ if one has both $\zeta \in O(\eta)$ and $\eta \in O(\zeta)$. Note that Θ is an equivalence relation, that is $\zeta \in \Theta(\eta) \Leftrightarrow \eta \in \Theta(\zeta)$.

Boshernitzan and Carroll give in [10] two upper bounds for $\rho^+(X)$ and $\sigma^+(X)$ for a subset X (Theorems 2.5 and 2.6 respectively) and a more precise estimation when the subset is a semi-interval (Theorem 2.8). Some slight modifications of the proofs can be made so that the results hold also for ρ^- and σ^- . We summarize these estimates in the following theorem.

Theorem 46 *For every $X \in \mathcal{A}([\ell, r[)$ one has $\rho^+(X), \rho^-(X) \in O(\Psi(X))$ and $\sigma^+(X), \sigma^-(X) \in O(1/|X|)$. Moreover, if T is minimal and J is a semi-interval, then $\rho^+(J) \in \Theta(\rho^-(J)) = \Theta(\sigma^+(J)) = \Theta(\sigma^-(J)) = \Theta(1/|J|)$.*

An immediate corollary of Theorem 46 is the following

Corollary 47 *Let T be minimal and assume that*

$$\{T^i(z) \mid -m+1 \leq i \leq n-1\} \cap J = \emptyset$$

for some point $z \in [\ell, r[$, some semi-interval $J \subset [\ell, r[$ and some integers $m, n \geq 1$. Then $|J| \in O(1/\max\{m, n\})$.

Proof. By the hypothesis, $z \notin \bigcup_{i=0}^{n-1} T^{-i}(J)$, then we have $\rho^-(J) \geq n$. By Theorem 46, we obtain $|J| \in \Theta(1/\rho^-(J)) \subseteq O(1/n)$. Symmetrically, since $\rho^+(J) \geq m$, one has $|J| \in O(1/m)$. Then $|J| \in O(\min\{1/m, 1/n\}) = O(1/\max\{m, n\})$. ■

4.3 Reduced complexity of admissible semi-intervals

In order to obtain Theorem 42, we prove some preliminary results concerning the reduced complexity of admissible semi-intervals.

Let T be an s -interval exchange transformation. Recall from Section 2 that we denote by $\text{Sep}(T) = \{\gamma_i \mid 0 \leq i \leq s-1\}$ the set of separation points. For every $n \geq 0$ define $\mathcal{S}_n(T) = \bigcup_{i=0}^{n-1} T^{-i}(\text{Sep}(T))$ with the convention $\mathcal{S}_0 = \emptyset$.

Since $\text{Sep}(T^{-1}) = T(\text{Sep}(T))$, one has $\mathcal{S}_n(T^{-1}) = T^{n-1}(\mathcal{S}_n(T))$.

Given two integers $m, n \geq 1$, we can define $\mathcal{S}_{m,n} = \mathcal{S}_m(T) \cup \mathcal{S}_n(T^{-1})$. An easy calculation shows that $\mathcal{S}_{m,n}(T) = \bigcup_{i=-m+1}^n T^i(\text{Sep}(T))$. Observe also that $\mathcal{S}_{m,n}(T) = T^n(\mathcal{S}_{m+n}(T)) = T^{-m+1}(\mathcal{S}_{m+n}(T))$.

Denote by $\mathcal{V}_{m,n}(T)$ the family of semi-intervals whose endpoints are in $\mathcal{S}_{m,n}(T)$. Put $\mathcal{V}(T) = \bigcup_{m,n \geq 0} \mathcal{V}_{m,n}(T)$. Every admissible semi-interval belongs to $\mathcal{V}(T)$, while the converse is not true.

Theorem 48 $\Pi(J) \in \Theta(1)$ *for every semi-interval J admissible for T .*

Proof. Let m, n be the two minimal integers such that $J = [t, w[\in \mathcal{V}_{m,n}(T)$. Then $t, w \in \{T^m(\gamma_i) \mid 1 \leq i \leq s\} \cup \{T^{-n}(\gamma_i) \mid 1 \leq i \leq s\}$. Suppose, for instance, $t = T^M(\gamma)$, with $M = \max\{m, n\}$ and γ a separation point. The other cases (namely, $t = T^{-M}(\gamma)$, $w = T^M(\gamma)$ or $w = T^{-M}(\gamma)$) are proved similarly.

The only semi-interval in $\mathcal{V}_{0,0}(T)$ is $[\ell, r[$ and clearly in this case the theorem is verified.

Suppose then that $J \in \mathcal{V}_{m,n}(T)$ for some nonnegative integers m, n with $m + n > 0$. We have $\Psi(J) = \max\{\Psi(t), \Psi(w)\} \leq Mu$ where u is the constant introduced in Proposition 44. Moreover, by the definition of admissibility one has $\{T^j(\gamma) \mid 1 \leq j \leq M\} \cap J = \emptyset$. Thus, by Corollary 47 we have $|J| \in O(1/M)$. Then $\Pi(J) = |J| \Psi(J) \in O(1)$. By Proposition 45 we have $\Pi(J) > 1/(4\sqrt{d})$. This concludes the proof. ■

Denote by $\mathcal{U}_{m,n}(T)$ the family of semi-intervals partitioned by $\mathcal{S}_{m,n}(T)$. Clearly $\mathcal{V}_{m,n}(T)$ contains $\mathcal{U}_{m,n}(T)$. Indeed every semi-interval $J \in \mathcal{V}_{m,n}(T)$ is a finite union of contiguous semi-intervals belonging to $\mathcal{U}_{m,n}(T)$.

Note that $\mathcal{U}_{m,0}(T)$ is the family of semi-intervals exchanged by T^m , while $\mathcal{U}_{0,n}(T)$ is the family of semi-intervals exchanged by T^{-n} .

Put $\mathcal{U}(T) = \bigcup_{m,n \geq 0} \mathcal{U}_{m,n}(T)$. Using Theorem 48 we easily deduce the following corollary, which is a generalization of Theorem 2.11 in [10].

Corollary 49 $\Pi(J) \in \Theta(1)$ for every semi-interval $J \in \mathcal{U}(T)$.

We are now able to prove Theorem 42.

Proof of Theorem 42. By Theorem 26, every admissible semi-interval can be obtained by a finite sequence χ of right and left Rauzy inductions. Thus we can enumerate the family of all admissible semi-intervals. The conclusion easily follows from Theorem 43 and Theorem 48. ■

An immediate corollary of Theorem 42 is the following.

Corollary 50 *Let T be a regular interval exchange transformation defined over a quadratic field. Then the induction graph $\mathcal{G}(T)$ and the modified induction graph $\tilde{\mathcal{G}}(T)$ are finite.*

Example 51 Let T be the regular interval exchange transformation of Example 8. The modified induction graph $\tilde{\mathcal{G}}(T)$ is represented in Figure 15. The transformation T belongs to the similarity class $\langle T_1 \rangle$ as well as transformations S of Example 31 and U of Example 33. The transformations $\psi(T)$ and $\psi^2(T)$ of Example 25 belongs respectively to classes $\langle T_2 \rangle$ and $\langle T_4 \rangle$, while the two last transformations of Figure 14, namely $\varphi\psi(T)$ and $\varphi^2\psi(T)$, belongs respectively to $\langle T_5 \rangle$ and $\langle T_7 \rangle$. Finally, the left Rauzy induction sequence from T to $U = \varphi^6(T)$ corresponds to the loop $\langle T_1 \rangle \rightarrow \langle T_3 \rangle \rightarrow \langle T_4 \rangle \rightarrow \langle T_6 \rangle \rightarrow \langle T_7 \rangle \rightarrow \langle T_8 \rangle \rightarrow \langle T_1 \rangle$ in $\tilde{\mathcal{G}}(T)$.

4.4 Primitive morphic sets

In this section we show an important property of interval exchange transformations defined over a quadratic field, namely that the related interval exchange sets are primitive morphic. Let prove first the following result.




Figure 15: Modified induction graph of the transformation T .

Proposition 52 *Let $T, \chi(T)$ be two equivalent regular interval exchange transformations with $\chi \in \{\varphi, \psi\}^*$. There exists a primitive morphism θ and a point $z \in D(T)$ such that the natural coding of T relative to z is a fixed point of θ .*

Proof. By Proposition 13, the set $F(T)$ is uniformly recurrent. Thus, there exists a positive integer N such that every letter of the alphabet appears in every word of length N of $F(T)$. Moreover, by Theorem 30, applying iteratively the Rauzy induction, the length of the domains tends to zero.

Consider $T' = \chi^m(T)$, for a positive integer m , such that $D(T') < \varepsilon$, where ε is the positive real number for which, by Lemma 11, the first return map for every point of the domain is “longer” than N , i.e. $T'(z) = T^{n(z)}(z)$, with $n(z) \geq N$, for every $z \in D(T')$.

By Theorem 38 and the remark following it, there exists an automorphism θ of the free group and a point $z \in D(T') \subseteq D(T)$ such that the natural coding of T relative to z is a fixed point of θ , that is $\Sigma_T(z) = \theta(\Sigma_T(z))$.

By the previous argument, the image of every letter by θ is longer than N , hence it contains every letter of the alphabet as a factor. Therefore, θ is a primitive morphism. ■

Theorem 53 *Let T be a regular interval exchange transformation defined over a quadratic field. The interval exchange set $F(T)$ is primitive morphic.*

Proof. By Theorem 42 there exists a regular interval transformation S such that we can find in the induction graph $\mathcal{G}(T)$ a path from $[T]$ to $[S]$ followed by a cycle on $[S]$. Thus, by Theorem 38 there exists a point $z \in D(S)$ and two

automorphisms θ, η of the free group such that $\Sigma_T(z) = \theta(\Sigma_S(z))$, with $\Sigma_S(z)$ a fixed point of η .

By Proposition 52 we can suppose, without loss of generality, that η is primitive. Therefore, $F(T)$ is a primitive morphic set. ■

Example 54 Let $T = T_{\lambda, \pi}$ be the transformation of Example 8 (see also 14). The set $F(T)$ is primitive morphic. Indeed the transformation T is regular and the length vector $\lambda = (1 - 2\alpha, \alpha, \alpha)$ belongs to $\mathbb{Q}[\sqrt{5}]^3$.

References

- [1] V. I. Arnol'd. Small denominators and problems of stability of motion in classical and celestial mechanics. *Uspehi Mat. Nauk*, 18(6 (114)):91–192, 1963.
- [2] Peter Balá z, Zuzana Masáková, and Edita Pelantová. Characterization of substitution invariant words coding exchange of three intervals. *Integers*, 8:A20, 21, 2008.
- [3] L'ubomíra Balková, Edita Pelantová, and Wolfgang Steiner. Sequences with constant number of return words. *Monatsh. Math.*, 155(3-4):251–263, 2008. (<http://dx.doi.org/10.1007/s00605-008-0001-2>).
- [4] Jean Berstel, Clelia De Felice, Dominique Perrin, Christophe Reutenauer, and Giuseppina Rindone. Bifix codes and Sturmian words. *J. Algebra*, 369:146–202, 2012. (<http://dx.doi.org/10.1016/j.jalgebra.2012.07.013>).
- [5] Valérie Berthé, Clelia De Felice, Francesco Dolce, Julien Leroy, Dominique Perrin, Christophe Reutenauer, and Giuseppina Rindone. Acyclic, connected and tree sets. *Monatsh. Math.*, 176(4):521–550, 2015. (<http://dx.doi.org/10.1007/s00605-014-0721-4>).
- [6] Valérie Berthé, Clelia De Felice, Francesco Dolce, Julien Leroy, Dominique Perrin, Christophe Reutenauer, and Giuseppina Rindone. Bifix codes and interval exchanges. *J. Pure Appl. Algebra*, 219(7):2781–2798, 2015. (<http://dx.doi.org/10.1016/j.jpaa.2014.09.028>).
- [7] Valérie Berthé, Clelia De Felice, Francesco Dolce, Julien Leroy, Dominique Perrin, Christophe Reutenauer, and Giuseppina Rindone. Maximal bifix decoding. *Discrete Math.*, 338(5):725–742, 2015. (<http://dx.doi.org/10.1016/j.disc.2014.12.010>).
- [8] Valérie Berthé and Michel Rigo. Preliminaries. In *Combinatorics, automata and number theory*, volume 135 of *Encyclopedia Math. Appl.*, pages 1–33. Cambridge Univ. Press, Cambridge, 2010.
- [9] Co Rentin Boissy and Erwan Lanneau. Dynamics and geometry of the Rauzy-Veech induction for quadratic differentials. *Ergodic Theory Dynam. Systems*, 29(3):767–816, 2009.
- [10] M. D. Boshernitzan and C. R. Carroll. An extension of Lagrange's theorem to interval exchange transformations over quadratic fields. *J. Anal. Math.*, 72:21–44, 1997. (<http://dx.doi.org/10.1007/BF02843152>).

- [11] Michael D. Boshernitzan. Rank two interval exchange transformations. *Ergodic Theory Dynam. Systems*, 8(3):379–394, 1988. (<http://dx.doi.org/10.1017/S0143385700004521>).
- [12] I. P. Cornfeld, S. V. Fomin, and Ya. G. Sinaĭ. *Ergodic theory*, volume 245 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, New York, 1982. Translated from the Russian by A. B. Sosinskiĭ.
- [13] Claude Danthony and Arnaldo Nogueira. Measured foliations on nonorientable surfaces. *Ann. Sci. École Norm. Sup. (4)*, 23(3):469–494, 1990.
- [14] Fabien Durand. A characterization of substitutive sequences using return words. *Discrete Math.*, 179(1-3):89–101, 1998. ([http://dx.doi.org/10.1016/S0012-365X\(97\)00029-0](http://dx.doi.org/10.1016/S0012-365X(97)00029-0)).
- [15] Sébastien Ferenczi and Luca Q. Zamboni. Languages of k -interval exchange transformations. *Bull. Lond. Math. Soc.*, 40(4):705–714, 2008. (<http://dx.doi.org/10.1112/blms/bdn051>).
- [16] N. Pytheas Fogg. *Substitutions in dynamics, arithmetics and combinatorics*, volume 1794 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2002. Edited by V. Berthé, S. Ferenczi, C. Mauduit and A. Siegel.
- [17] Yann Jullian. An algorithm to identify automorphisms which arise from self-induced interval exchange transformations. *Math. Z.*, 274(1-2):33–55, 2013. (<http://dx.doi.org/10.1007/s00209-012-1056-2>).
- [18] Jacques Justin and Laurent Vuillon. Return words in Sturmian and episturmian words. *Theor. Inform. Appl.*, 34(5):343–356, 2000. (<http://dx.doi.org/10.1051/ita:2000121>).
- [19] Michael Keane. Interval exchange transformations. *Math. Z.*, 141:25–31, 1975.
- [20] Petr Kůrka. *Topological and symbolic dynamics*, volume 11 of *Cours Spécialisés [Specialized Courses]*. Société Mathématique de France, Paris, 2003.
- [21] M. Lothaire. *Algebraic combinatorics on words*, volume 90 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2002. A collective work by Jean Berstel, Dominique Perrin, Patrice Seebold, Julien Cassaigne, Aldo De Luca, Steffano Varricchio, Alain Lascaux, Bernard Leclerc, Jean-Yves Thibon, Veronique Bruyere, Christiane Frougny, Filippo Mignosi, Antonio Restivo, Christophe Reutenauer, Dominique Foata, Guo-Niu Han, Jacques Desarmenien, Volker Diekert, Tero Harju, Juhani Karhumaki and Wojciech Plandowski, With a preface by Berstel and Perrin.
- [22] Tomasz Miernowski and Arnaldo Nogueira. Exactness of the Euclidean algorithm and of the Rauzy induction on the space of interval exchange transformations. *Ergodic Theory Dynam. Systems*, 33(1):221–246, 2013. (<http://dx.doi.org/10.1017/S014338571100085X>).
- [23] V. I. Oseledec. The spectrum of ergodic automorphisms. *Dokl. Akad. Nauk SSSR*, 168:1009–1011, 1966.
- [24] Gérard Rauzy. Échanges d’intervalles et transformations induites. *Acta Arith.*, 34(4):315–328, 1979.

- [25] Alexandra Skripchenko. Symmetric interval identification systems of order three. *Discrete Contin. Dyn. Syst.*, 32(2):643–656, 2012. (<http://dx.doi.org/10.3934/dcds.2012.32.643>).
- [26] Marcelo Viana. Ergodic theory of interval exchange maps. *Rev. Mat. Complut.*, 19(1):7–100, 2006.
- [27] Laurent Vuillon. On the number of return words in infinite words constructed by interval exchange transformations. *Pure Math. Appl. (P.U.M.A.)*, 18(3-4):345–355, 2007.
- [28] Jean-Christophe Yoccoz. Interval exchange maps and translation surfaces. In *Homogeneous flows, moduli spaces and arithmetic*, volume 10 of *Clay Math. Proc.*, pages 1–69. Amer. Math. Soc., Providence, RI, 2010.